

Activités physiques et santé

Bibliographie complète

Patrick Laure

© Éditions Ellipses, 2007

Collection « L'essentiel en sciences du sport »

Certaines références sont citées à plusieurs reprises dans le texte. Elles n'apparaissent qu'une seule fois ici, dans le chapitre où elles sont mentionnées en premier.

Généralités

Abu-Omar K, Rütten A, Robine J.-M. Self-rated health and physical activity in the European Union. *Soz Präventivmed* 2004 ; 49 : 235-42.

Annual Global Move for Health Initiative. A Concept Paper. World Health Organization. WHO/NMH/NPH/PAH/03.1

Bernstein M, Morabia A, Sloutskis D. Definition and prevalence of sedentarism in an urban population. *Am J Public Health* 1999 ; 89 : 862-67.

Blair SN, Cheng Y, Holder JS. Is physical activity or physical fitness more important in defining health benefits ? *Med Sci Sports Exerc* 2001 ; 33 : 379-99.

Bouchard C, Shephard RJ. Physical activity, fitness, and health : the model and the key concepts. In : Bouchard C, Shephard RJ, Stephens T, eds. *Physical activity, fitness and health*. Champaign, IL : Human Kinetics, 1994 : 77-88.

Bouchard C. Individual differences in the response to regular exercise. *Int J Obes Relat Metab Disord* 1995 ; 19 : S5-8.

Centers for Disease Control and Prevention (CDC). Trends in strength training-United States, 1998-2004. *Morb Mortal Wkly Rep* 2006 ; 55 : 769-72.

Centers for Disease Control and Prevention. *DATA2010, the Healthy People 2010 Database*. July, 2004.

Charte d'Ottawa pour la promotion de la santé. WHO/HPR/HEP/95.1. OMS, Genève, 1986.

Colditz GA. Economic costs of obesity and inactivity. *Med Sci Sports Exerc* 1999 ; 31 : S663-7.

Craig, CL. *Physical Activity and the Renewal of the Health Care System*. Ottawa : Institut canadien de la recherche sur la condition physique et le mode de vie, 1994.

FAO/WHO/UNU. Energy and protein requirements. Report of a joint FAO/WHO/UNU expert consultation. *World Health Organ Tech Rep Ser* 1985 ; 724 : 34-52.

Garrett NA, Brasure M, Schmitz KH, et al. Physical inactivity : direct cost to a health plan. *Am J Prev Med* 2004 ; 27 : 304-9.

Goetzl RZ, Anderson DR, Whitmer RW, et al. The relationship between modifiable health risks and health care expenditures : an analysis of the multi-employer HERO health risk and cost database. The Health Enhancement Research Organization (HERO) Research Committee. *J Occup Environ Med* 1998 ; 40 : 843-54.

Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ) - Short Form. Version 2.0. April 2004. www.ipaq.ki.se.

Guy WA. Contribution to a knowledge of the influence of employments upon health. *J Roy Stat Soc* 1843 ; 6 : 197-211.

Ham SA, Yore MM, JE Fulton JE, Kohl HW. Prevalence of No Leisure-Time Physical Activity -35 States and the District of Columbia, 1988-2002. *MMWR Morb Mortal Wkly Rep* 2004 ; 53 : 82-6.

Hardman AE, Stensel DJ. *Physical activity and health. The evidence explained*. London : Routledge, 2003.

Haut Comité de la santé publique. *Pour une politique nutritionnelle de santé publique en France : enjeux et recommandations*. Rapport au Secrétaire d'État à la Santé et aux Handicapés, Paris, juin 2000.

- Hillsdon M, Foster C, Thorogood M. Interventions for promoting physical activity. *The Cochrane Database of Systematic Reviews* 2005, Issue 1. Art. No : CD003180.pub2. DOI : 10.1002/14651858.CD003180.pub2.
- Katzmarzyk PT, Gledhill N, Shephard RJ. The economic burden of physical inactivity in Canada. *CMAJ* 2000 ; 163 : 1435-40.
- Katzmarzyk PT, Janssen I. The economic costs associated with physical inactivity and obesity in Canada : an update. *Can J Appl Physiol* 2004 ; 29 : 90-115.
- Keeler EB, Manning WG, Newhouse JP, et al. The external costs of a sedentary life-style. *Am J Public Health* 1989 ; 79 : 975-81.
- Koufaki P, Nash PF, Mercer TH. Assessing the efficacy of exercise training in patients with chronic disease. *Med Sci Sports Exerc* 2002 ; 34 : 1234-41.
- Leon AS, Gaskill SE, Rice T, et al. Variability in the response of HDL cholesterol to exercise training in the HERITAGE Family Study. *Int J Sports Med* 2002 ; 23 : 1-9.
- Martin BW, Beeler I, Szucs T, et al. Répercussions économiques du rapport entre santé et activité physique : premières estimations pour la Suisse. *Schweiz Z Sportmed Sporttraumatol* 2001 ; 49 : 87-9.
- Martinson BC, Crain AL, Pronk NP, et al. Changes in physical activity and short-term changes in health care charges : a prospective cohort study of older adults. *Prev Med* 2003 ; 37 : 319-26.
- Morris JN, Heady JA, Raffle PA, et al. Coronary heart disease and physical activity of work. *Lancet* 1953 ; 1053-1057 : 1111-20.
- Pronk NP, Goodman MJ, O'Connor PJ, Martinson BC. Relationship between modifiable health risks and short-term health care charges. *JAMA* 1999 ; 282 : 2235-9.
- Santé Canada. *Le fardeau économique de la maladie au Canada, 1993*. Ottawa : Ministre des Travaux publics et des services gouvernementaux du Canada, 1997.
- Smala A, Beeler I, Szucs T. *Die Kosten der körperlichen Inaktivität in der Schweiz*. Zürich : Abteilung für medizinische Ökonomie des Instituts für Sozial- und Präventivmedizin und des Universitätsspitals, 2001.
- Temple VA, Frey GC, Stanish HI. Physical activity of adults with mental retardation : review and research needs. *Am J Health Promot* 2006 ; 21 (1) : 2-12.
- US Department of Health and Human Services. *Healthy People 2010*. Washington DC, Conference Edition, vol. 2 ; 2000.
- Varo JJ, Martínez-González MA, de Irala-Estévez J, et al. Distribution and determinants of sedentary lifestyles European Union. *Int J Epidemiology* 2003 ; 32 : 138-46.
- Wang G, Pratt M, Macera CA, et al. Physical activity, cardiovascular disease, and medical expenditures in U.S. adults. *Ann Behav Med* 2004 ; 28 : 88-94.



Évaluation de l'activité physique d'une personne

MÉTHODES GÉNÉRALES D'ÉVALUATION

- Aadahl M, Jorgensen T. Validation of a new self-report instrument for measuring physical activity. *Med Sci Sports Exerc* 2003 ; 35 : 1196-202.
- Acheson KJ, Campbell IT, Edholm OG, et al. The measurement of daily energy expenditure : an evaluation of some techniques. *Am J Clin Nutr* 1980 ; 33 : 1155-64.
- Ainsworth BE, Haskell WL, Whitt MC, et al. Compendium of Physical Activities : An update of activity codes and MET intensities. *Med Sci Sports Exerc* 2000 ; 32 : S498-S516.
- Ainsworth BE, Jacobs DR, Leon AS, et al. Assessment of the accuracy of physical activity questionnaire occupational data. *J Occupational Med* 1993 35 : 1017-27.
- Baecke JAH, Burema J, Ftijters JER. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* 1982 ; 36 : 936-42.
- Bennett T, Campagna P. The utility of the CSA accelerometer to assess daily step count activity. *Can J Appl Physiol* 2002 ; 27 (S5) : np.

Berthouze SE, Minaire PM, Chatard JC, et al. A new tool for evaluating energy expenditure : the « QAPSE » development and validation. *Med Sci Sports Exerc* 1993 ; 25 : 1405-14.

Berthouze-Aranda SE, Aranda P. Un logiciel pour la mesure de l'activité physique : le PAQAP. *Sci Sports* 2001 ; 16 : 171-2.

Black AE, Cole TJ. Within- and between-subject variation in energy expenditure as measured by the doubly-labelled water technique : implications for validating reported dietary energy intake. *Eur J Clin Nutr* 2000 ; 54 : 386-94.

Blair SN, Haskell WL, Ho P, et al. Assessment of habitual physical activity by a seven-day recall in a community survey and controlled experiments. *Am J Epidemiol* 1985 ; 122 : 794-804.

Bouchard C, Tremblay A, Leblanc C, et al. A method to assess energy expenditure in children and adults. *Am J Clin Nutr* 1983 ; 37 : 461-7.

Bowles HR, FitzGerald SJ, Morrow JR, et al. Construct Validity of Self-reported Historical Physical Activity. *Am J Epidemiol* 2004 160 : 279-86.

Brage S, Wedderkopp N, Frankes PW, Andersen LB, Froberg K. Reexamination of validity and reliability of the CSA monitor in walking and running. *Med Sci Sports Exerc* 2003 ; 8 : 1447-54.

Brooks AG, Gunn SM, Withers RT, et al. Predicting Walking MET and Energy Expenditure from Speed or Accelerometry. *Med Sci Sports Exerc* 2005 ; 37 : 1216-23.

Bruce RA, Fisher LD, Cooper MN, Grey GO. Separation of effects of cardiovascular disease and age on ventricular function with maximal exercise. *Am J Cardiol* 1974 ; 34 : 757-763.

Busse ME, Wiles CM, van Deursen RMW. Community walking activity in neurological disorders with leg weakness. *J Neurol Neurosurg Psychiatry* 2006 ; 77 : 359-62.

Caspersen CJ, Blaemberg PM, Saris WHM, et al. The prevalence of selected physical activities and their relation with coronary heart disease risk factors in elderly men : the Zutphen study, 1985. *Am J Epidemiol* 1991 ; 133 : 1078-91.

CDC. Prevalence of Physical Activity, Including Lifestyle Activities Among Adults-United States, 2000-2001. *Morbidity and Mortality Weekly Report* 2003 ; 52 : 764-9.

Chasan-Taber L, Schmidt MD, Roberts DE, et al. Development and validation of a pregnancy physical activity questionnaire. *Med Sci Sports Exerc* 2004 ; 36 : 1750-60.

Coleman KJ, Epstein LH. Application of generalizability theory to measurement of activity in males who are not regularly active : a preliminary report. *Res Q Exerc Sport* 1998 ; 69 : 58-63.

Conway JM, Seale JL, Jacobs DR Jr, et al. Comparison of energy expenditure estimates from doubly labeled water, a physical activity questionnaire, and physical activity records. *Am J Clin Nutr* 2002 ; 75 : 519-25.

Craig CL, Marshall AL, Sjostrom M, et al. International physical activity questionnaire : 12-country reliability and validity. *Med Sci Sports Exerc* 2003 ; 35 : 1381-95.

Craig CL, Russell SJ, Cameron C. Reliability and validity of Canada's Physical Activity Monitor for assessing trends. *Med Sci Sports Exerc* 2002 ; 34 : 1462-7.

Dan AJ, Wilbur JE, Hedricks C, et al. Lifelong physical activity and older women. *Psychol Women Quarterly* 1990 ; 14 : 531-42.

Dehart-Beberley M, Foster C, Porcari JP, et al. Relationship between the talk test and the ventilatory threshold. *Clin Exerc Physiol* 2000 ; 2 : 34-8.

DiPietro L, Caspersen CJ, Ostfeld AM, Nadel ER. A survey for assessing physical activity among older adults. *Med Sci Sports Exerc* 1993 ; 25 : 628-42.

Dominguez-Berjon F, Borrell C, Nebot M, Plansencia A. Physical activity assessment in population surveys : can it really be simplified ? *Int J Epidemiol* 1999 ; 28:23-7.

Duncan GE, Sydemann SJ, Perri MG, et al. Can sedentary adults accurately recall the intensity of their physical activity ? *Prev Med* 2001 ; 33 : 18-26.

Duncan M, Black A, Yewdale J, Schofield G, Mummery K. Accuracy of two makes of electronic pedometers for measuring step counts at various speeds. *Australian conference of science and medicine in sport 2002 : Sports medicine and science at the extremes*. 12-16 October 2002, Melbourne.

Edwards JE, Babington P, Ozmun JC, et al. Validity and reliability of an activity monitor with the capability of providing geographic location using GPS. *Med Sci Sports Exerc* 2002 ; 5 : S41.

Fogelholm M, Malmberg J, Suni J, et coll. International Physical Activity Questionnaire : Validity against fitness. *Med Sci Sports Exerc* 2006 ; 38 : 753-60.

Foster RC, Lanningham-Foster LM, Manohar C, et al. Precision and accuracy of an ankle-worn accelerometer-based pedometer in step counting and energy expenditure. *Prev Med* 2005 ; 41 : 778-83.

Fox SM 3rd, Naughton JP, Haskell WL. Physical activity and the prevention of coronary heart disease. *Ann Clin Res* 1971 ; 3 : 404-32.

Freedson PS, Melanson EL, Sirard JR. Calibration of the Computer Science and Applications, Inc. accelerometer. *Med Sci Sports Exerc* 1998 ;30 : 777-81.

Garet M, Degache F, Costes F, et al. DAQIHF : methodology and validation of a daily activity questionnaire in heart failure. *Med Sci Sports Exerc* 2004 ; 36 : 1275-82.

Ginis KA, Latimer AE, Hicks AL, Craven BC. Development and evaluation of an activity measure for people with spinal cord injury. *Med Sci Sports Exerc* 2005 ; 37 : 1099-111.

Godin G, Shephard RJ. A simple method to assess exercise behavior in the community. *Can J Appl Sport Sci* 1985 10 : 14146.

Guilbert P, Baudier F, Gautier A (dir). *Baromètre Santé 2000. Vol 2 : Résultats*. Vanves : CFES, 2001.

Haskell WL, Yee MC, Evans A, Irby PJ. Simultaneous measurement of heart rate and body motion to quantitate physical activity. *Med Sci Sports Exerc* 1993 ; 25 : 109-15.

Haskell WL, Taylor HL, Wood PD, et al. Strenuous physical activity, treadmill exercise test performance and plasma high-density lipoprotein cholesterol. *Circulation* 1980 ; 62 (Suppl. IV) : 53-61.

Health Canada. *Physical activity of Canadians. 2.10 Non-leisure physical activity*. National Population Health Survey Highlights, November 1999 ; 2.

Hendelman D, Miller K, Baggett C, et al. Validity of accelerometry for the assessment of moderate intensity physical activity in the field. *Med Sci Sports Exerc* 2000 ; 32 : S442-9.

Hopkins WG, Wilson NC, Russell DG. Validation of the physical activity instrument for the life in New Zealand national survey. *Am J Epidemiol* 1991 ; 133 : 73-82.

Hornbuckle LM, Bassett DR, Thompson DL. Pedometer-Determined Walking and Body Composition Variables in African-American Women. *Med Sci Sports Exerc* 2005 ; 37 : 1069-74.

Hultquist CN, Albright C, Thompson DL. Comparison of Walking Recommendations in Previously Inactive Women. *Med Sci Sports Exerc* 2005 ; 37 : 676-83.

Inbar O, Oren A, Scheinowitz M, et al. Normal cardiopulmonary responses during incremental exercise in 20-70-yr-old men. *Med Sci Sport Exerc* 1994 ;26 : 538-46.

Irwin ML, Ainsworth BE, Conway JM. Estimation of Energy Expenditure from Physical Activity Measures : Determinants of Accuracy. *Obes Res* 2001 ; 9 : 517-25.

Jacobs D, Ainsworth B, Hartman T, Leon A. A simultaneous evaluation of 10 commonly used physical activity questionnaires. *Med Sci Sports Exerc* 1993 ; 25 : 81-91.

Jacobs DR, Hahn L, Haskell WL et al. Validity and reliability of a short physical activity history : CARDIA and the Minnesota Heart Health Program. *J Cardiopulm Rehab* 1989 ; 9 : 448-59.

Jacobs DR, Luepker RY, Mittelmark MB et al. Community-wide prevention strategies : evaluation design of the Minnesota Heart Health Program. *J Chronic Dis* 1986 ; 39 : 775-88.

Kaleta D, Jegier A. Occupational energy expenditure and leisure-time physical activity. *Int J Occup Med Environ Health* 2005 ; 18 : 351-6.

Kannel WB, Sorlie P. Some health benefits of physical activity. The Framingham study. *Arch Intern Med* 1979 ; 139 : 857-61.

Klute GK, Berge JS, Orendurff MS, et al. Prosthetic intervention effects on activity of lower-extremity amputees. *Arch Phys Med Rehabil* 2006 ; 87 : 717-22.

Kohl HW, Blair SN, Paffenbarger RS Jr, et al. A mail survey of physical activity habits as related to measured physical fitness. *Am J Epidemiol* 1988 ; 127 : 1228-39.

Kriska A. Ethnic and cultural issues in assessing physical activity. *Res Q* 2000 ; 71 : S47-53.

Kriska AM, Black-Sandler R, Cauley JA, et al. The assessment of historical physical activity and its relation to adult bone parameters. *Am J Epidemiol* 1988 127 : 1053-63.

Kriska AM, Knowler WC, LaPorte RE et al. Development of a questionnaire to examine relationship of physical activity and diabetes in Pima Indians. *Diabetes Care* 1990 ; 13 : 401-11.

La pratique des activités physiques et sportives en France. Paris : Ministère des sports et INSEP, 2005.

Lamb KL, Brodie DA. Leisure-time physical activity as an estimate of physical fitness : a validation study. *J Clin Epidemiol* 1991 ;44 : 41-52.

Lamonte MJ, Ainsworth BE. Quantifying energy expenditure and physical activity in the context of dose response. *Med Sci Sports Exerc* 2001 ; 33 : S370-8.

Larsson P. Global positioning system and sport-specific testing. *Sports Med* 2003 ; 33 : 1093-101.

Leenders NY, Neslon TE, Sherman WM. Ability of different physical activity monitors to detect movement during treadmill walking. *Int J Sports Med* 2003 ; 24 : 43-50

Leon AS, Jacob DR, DeBacker G, Taylor HL. Relationship of physical characteristics and life habits to treadmill exercise capacity. *Am J Epidemiol* 1981 : 113 : 653-60.

Lester M, Sheffield LT, Trammel P, Reeves TJ. The effect of age and athletic training on the maximal heart rate during muscular exercise. *Am Heart J* 1968 ; 76 : 370-6.

Levin S, Jacobs DR, Ainsworth BE, et al. Intra-individual variation and estimates of usual activity. *Ann Epidemiol* 1999 ; 9 : 481-8.

Lof M, Hannestad U, Forsum E. Comparison of commonly used procedures, including the doubly-labelled water technique, in the estimation of total energy expenditure of women with special reference to the significance of body fatness. *Br J Nutr* 2003 ; 90 : 961-8

Londeree BR, Moeschberger ML. Effect of age and other factors on maximal heart rate. *Res Quarter Exerc Sport* 1982 ; 53 : 297-304.

Lowther MC, Mutrie N, Loughlan C, McFarlane C. Development of a Scottish physical activity questionnaire : a tool for use in physical activity interventions *Br J Sports Med* 1999 ; 33 : 244-9.

Magnus K, Matroos A, Strackee J. Walking, cycling, or gardening, with or without seasonal interruption. in relation to acute coronary events. *Am J Epidemiol* 1979 ; 110 : 724-33.

Marshall AL, Smith BJ, Bauman AE, Kaur S. Reliability and validity of a brief physical activity assessment for use by family doctors. *Br J Sports Med* 2005 ; 39 : 294-7.

Mâsse LC, Fulton JE, Watson KL, et al. Influence of body composition on physical activity validation studies using doubly labeled water. *J Appl Physiol* 2004 ; 96 : 1357-64.

Mathews JE, Freedson PS. Field trial of a three-dimensional activity monitor : comparison with self report. *Med Sci Sports Exerc* 1995 ; 27 : 1071-8.

Matthews CE, Ainsworth BE, Hanby C, et al. Development and Testing of a Short Physical Activity Recall Questionnaire. *Med Sci Sports Exerc* 2005 ; 37 : 986-94.

Matthews CE, Ainsworth BE, Thompson RW, Bassett DR Jr. Sources of variance in daily physical activity levels as measured by an accelerometer. *Med Sci Sports Exerc* 2002 ; 34 : 1376-81.

Matthews CE, Shu XO, Yang G, et al. Reproducibility and validity of the Shanghai Women's Health Study physical activity questionnaire. *Am J Epidemiol* 2003 ; 158 : 1114-22.

McCormack G, Giles-Corti B, Milligan R. Demographic and individual correlates of achieving 10,000 steps/day : use of pedometers in a population-based study. *Health Promot J Austr* 2006 ; 17 : 43-7.

McDonald CM, Widman LM, Walsh DD, et al. Use of step activity monitoring for continuous physical activity assessment in boys with Duchenne muscular dystrophy. *Arch Phys Med Rehabil* 2005 ; 86 : 802-8.

Miller DJ, Freedson PS, Kline GM. Comparison of activity levels using the Caltrac accelerometer and five questionnaires. *Med Sci Sports Exerc* 1994 ; 26 : 376-82.

Morris JN, Adasn C, Chave SPW, Sirey C. Vigorous exercise in leisure-time and the incidence of coronary heart disease. *Lancet* 1973 ; 1 : 333-39.

Morrison JF, Van Malsen S, Noakes TD. Leisure-time physical activity levels, cardiovascular fitness and coronary risk factors in 1015 white Zimbabweans. *South Africa Med J* 1984 ; 65 : 250-56.

Motl RW, McAuley E, DiStefano C. Is social desirability associated with self-reported physical activity ? *Prev Med* 2005 ; 40 : 735-9.

Paffenbarger RS Jr, Blair SN, Lee IM, Hyde RT. Measurement of physical activity to assess health effects in free-living populations. *Med Sci Sports Exerc* 1993 ; 25 : 60-70.

Paffenbarger RS Jr, Wing AL, Hyde RT. Physical activity as an index of heart attack risk in college alumni. *Am J Epidemiol* 1978 ; 108 : 161-75.

- Parker DL, Leaf DA, Mc Mee SR. Validation of a new questionnaire for the assessment of leisure-time physical activity. *Ann Sports Med* 1988 ; 4 : 72-81.
- Persinger R, Foster C, Gibson M, et al. Consistency of the Talk Test for Exercise Prescription. *Med Sci Sports Exerc* 2004 ; 36 : 1632-6.
- Pollock ML. The quantification of endurance training programs. *Exerc Sport Sci Rev* 1973 ; 1 : 155-88.
- Pratiques et connaissances des Réunionnais vis-à-vis du diabète, de l'alimentation et de l'activité physique.* Saint-Denis, ORS : 2004.
- Prentice A. Stable isotopes in nutritional sciences and the study of energy metabolism. *Scand J Nutr* 1999 ;43 : 56-62.
- Ravussin E, Harper IT, Rising R, Bogardus C. Energy expenditure by doubly labeled water : validation in lean and obese subjects. *Am J Physiol* 1991 ; 261 : E402-9.
- Recalde PT, Foster C, Skempt-Arlt KM, et al. The talk test as a simple marker of ventilatory threshold. *S Afr J Sports Med* 2002 ; 8 : 5-8.
- Reiff GG, Montoye HJ, Remington RD, et al. Assessment of physical activity by questionnaire and interview. *J Sports Med Phys Fitness* 1967 ; 7 : 135-42.
- Rennie K, Wareham N. The validation of physical activity instruments for measuring energy expenditure : problems and pit falls. *Public Health Nutr* 1998 ; 1 : 265-71.
- Robergs RA, Landwehr R. The surprising history of the « HRmax = 220-age » equation. *JEPonline* 2002 ; 5 : 1-10.
- Robinson S. Experimental studies of physical fitness in relation to age. *Arbeitsphysiologie* 1938 ; 10 : 251-323.
- Rosenberg M, Bull FC, Braham R, et al. Assessing the accumulation of moderate-intensity physical activity in adults. *Med Sci Sports Exerc* 2002 ; 5 : S207.
- Rowlands AV, Eston RG, Ingledew DK. Relationship between activity levels, aerobic fitness, and body fat in 8- to 10-yr-old children. *J Appl Physiol* 1999 ; 86 : 1428-35.
- Sallis JF, Haskell WL, Wood PD et al. Physical activity assessment methodology in the five-city project. *Am J Epidemiol* 1985 ; 121 : 91-106.
- Saltin B, Grimby G. Physiological analysis of middle-aged and old former athletes. *Circulation* 1968 ; 38 : 1104-15.
- Schmidt MD, Freedson PS, Chasan-Taber L. Estimating physical activity using the CSA accelerometer and a physical activity log. *Med Sci Sports Exerc* 2003 ; 35 : 1615-11.
- Schneider PL, Crouter SE, Bassett DR. Pedometer measures of free-living physical activity : comparison of 13 models. *Med Sci Sports Exerc* 2004 ; 2 : 331-5.
- Schneider PL, Crouter SE, Lukajic O Bassett DR. Accuracy and Reliability of 10 Pedometers for Measuring Steps over a 400-m Walk. *Med Sci Sports Exerc.*2003 ;10,1779-84.
- Schutz Y, Chambaz A. Could a satellite-based navigation system (GPS) be used to assess the physical activity of individuals on earth ? *Eur J Clin Nutr* 1997 ; 51 : 338-9.
- Schutz Y, Herren R. Assessment of speed of human locomotion using a differential satellite global positioning system. *Med Sci Sports Exerc* 2000 ; 32 : 642-6.
- Shapiro S, Wemblart E, Frank CW, Sager RV. The HIP study of incidence and prognosis of coronary heart disease. *J Chron Dis* 1965 ; 18 : 527-58.
- Shepard RJ. Limits to the measurement of habitual physical activity by questionnaires. *Br J Sports Med* 2003 ; 37 : 197-206
- Shepherd EF, Toloza E, McClung CD, Schmalzried TP. Step activity monitor increased accuracy in quantifying ambulatory activity. *J Orthop Res* 1999 ; 17 : 703-8.
- Sirard JR, Melanson EL, Freedson PS. Field evaluation of the Computer Science and Applications, Inc. physical activity monitor. *Med. Sci. Sports Exerc* 2000 ;3 : 695-700.
- Snitker S, Tataranni PA, Ravussin E. Spontaneous physical activity in a respiratory chamber is correlated to habitual physical activity. *Int J Obes Relat Metab Disord* 2001 ; 25 : 1481-6.
- Sobngwi E, Mbanya JC, Unwin NC, et al. Development and validation of a questionnaire for the assessment of physical activity in epidemiological studies in Sub-Saharan Africa. *Int J Epidemiol* 2001 ; 30 : 1361-8.
- Song KM, Bjornson KF, Cappello T, Coleman K. Use of the StepWatch activity monitor for characterization of normal activity levels of children. *J Pediatr Orthop* 2006 ; 26 : 245-9.
- Speakman J. The history and theory of the doubly labeled water technique. *Am J Clin Nutr* 1998 ; 68 : S932-S8.

Starling RD, Matthews DE, Ades PA, Poehlman ET. Assessment of physical activity in older individuals : a doubly labeled water study. *J Appl Physiol* 1999 ; 86 : 2090-6.

Steffen LM, Arnett DK, Blackburn H, et al. Population trends in leisure-time physical activity : Minnesota Heart Survey, 1980-2000. *Med Sci Sports Exerc* 2006 ; 38 : 1716-23.

Steffen-Batey L, Nichaman MZ, Goff DC, et al. Change in Level of Physical Activity and Risk of All-Cause Mortality or Reinfarction : The Corpus Christi Heart Project. *Circulation* 2000 ; 102 : 2204-9.

Sunnegardh J, Bratteby L, Sjolin S. Physical activity and sports involvement of 8- and 13-year-old children in Sweden. *Acta Paediatr Scand* 1985 ; 74 : 904-12.

Suzuki I, Kawakami N, Shimizu H. Reliability and validity of a questionnaire for assessment of energy expenditure and physical activity in epidemiological studies. *J Epidemiol* 1998 ; 8 : 152-9.

Taylor HL, Jacobs DR Jr, Schecker B, et al. A questionnaire for the assessment of leisure-time physical activities. *J Chron Dis* 1978 ; 31 : 741-55.

Torgen M, Alfredsson L, Koster M, et al. Reproducibility of a questionnaire for assessment of present and past physical activities. *Int Arch Occup Environ Health* 1997 ; 70 : 107-18.

Tudor-Locke C, Bassett DR. How many steps/day are enough ? Preliminary pedometer indices for public health. *Sports Medicine* 2004 ; 34 : 1-8.

Tudor-Locke C, Bassett J, Swartz AM, et al. A preliminary study of one year of pedometer self-monitoring. *Ann Behav Med* 2004 ; 28 : 158-62.

Tudor-Locke C, Burkett L, Reis JP, et al. How many days of pedometer monitoring predict weekly physical activity in adults ? *Prev Med* 2005 ; 40 : 293-8.

Tudor-Locke C, Myers AM, Bell RC, et al. Preliminary outcome evaluation of The First Step Program : a daily physical activity intervention for individuals with type 2 diabetes. *Patient Educ Couns* 2002 ; 47 : 23-8.

Tudor-Locke C, Sisson SB, Collova T, et al. Pedometer-determined step count guidelines for classifying walking intensity in a young ostensibly healthy population. *Can J Appl Physiol* 2005 ; 30 : 666-76.

Tudor-Locke C, Williams JE, Reis JP, Pluto D. Utility of pedometers for assessing physical activity : construct validity. *Sports Med* 2004 ; 34 : 281-91.

Van Aalst, Kazakov D, McLean G. *SPARC Facts 1997-1998-2000*. Wellington : 2002.

van Sluijs EM, van Poppel MN, Twisk JW, van Mechelen W. Physical activity measurements affected participants' behavior in a randomized controlled trial. *J Clin Epidemiol* 2006 ; 59 : 404-11.

Verschuur R, Kemper H. Habitual physical activity. *Med Sci Sports Exerc* 1985 ; 20 : 56-65.

Vincent SD, Pangrazi RP. An examination of the activity patterns of elementary school children. *Pediatr Exerc Sci* 2002 ; 14 : 432-41

Voorrips LE, Ravelli ACI, Dongelmans PCA, et al. A physical activity questionnaire for the elderly. *Med Sci Sports Exerc* 1991 ; 23 : 974-79.

Vuillemin A, Guillemin F, Denis G, et al. A computer-assisted assessment of lifetime physical activity : reliability and validity of the QUANTAP software. *Rev Epidemiol Santé Pub* 2000 ; 48 : 157-67.

Vuillemin A, Oppert JM, Guillemin F, et al. Self-administered questionnaire compared with interview to assess past-year physical activity. *Med Sci Sports Exerc* 2000 ; 32 : 1119-24.

Walsh MC, Gary R, Hunter GR, et al. Comparison of self-reported with objectively assessed energy expenditure in black and white women before and after weight loss. *Am J Clin Nutr* 2004 ; 79 : 1013-9.

Warms C. Physical activity measurement in persons with chronic and disabling conditions : methods, strategies, and issues, *Fam Comm Health* 2006 ; 29 : S78-88S.

Washburn RA, Adams JL, Haile GT. Physical activity assessment for epidemiologic research : the utility of two simplified approach. *Prev Med* 1987 ; 16 : 636-46.

Washburn RA, Goldfield SR, Smith KW, McKinlay JB. The validity of self-reported exercise-induced sweating as a measure of physical activity. *Am J Epidemiol* 1990 ; 132 : 107-13.

Washburn RA, Jacobsen DJ, Sonko BJ, et al. The validity of the Stanford Seven-Day Physical Activity Recall in young adults. *Med Sci Sports Exerc* 2003 ; 8 : 1374-80.

Weinsier RL, Hunter GR, Zuckerman PA, et al. Energy expenditure and free-living physical activity in black and white women : comparison before and after weight loss. *Am J Clin Nutr* 2000 ; 71 : 1138-46.

Welk GJ, Differding JA, Thompson RW et al. The utility of the Digi-walker step counter to assess daily physical activity patterns. *Med Sci Sports Exerc* 2000 ; 32 : S481-S8.

- Wendel-Vos GC, Schuit AJ, Saris WH, Kromhout D. Reproducibility and relative validity of the Short Questionnaire to Assess Health-enhancing physical activity. *J Clin Epidemiol* 2003 ; 56 : 1163-9.
- Westerterp K. Limits to sustainable human metabolic rate. *J Experiment Biol* 2001 ; 204 : 3183-7.
- Westerterp KR. Physical activity assessment with accelerometers. *Int J Obes Relat Metab Disord* 1999 ; 23 : S45-9.
- Wilcox S, Irwi ML, Addy C, et al. Agreement between participant-rated and compendium-coded intensity of daily activities in a triethnic sample of women ages 40 years and older. *Ann Behav Med* 2001 ; 23 : 253-6.
- Wilde BE, Sidman CL, Corbin CB. A 10000 step count as a physical activity target for sedentary women. *Res Q Exerc Sport* 2001 ; 72 : 411-4.
- Winters-Hart CS, Brach JS, Storti KL, et al. Validity of a questionnaire to assess historical physical activity in older women. *Med Sci Sports Exerc* 2004 ; 36 : 2082-7.
- World Health Organization. *Energy and Protein : Requirements. Report of a Joint FAO/WHO/UNU Expert Consultation*. Technical Report Series no. 724. Geneva : WHIO, 1985.
- Yasin S, Alderson MR, Marr JW, et al. Assessment of habitual physical activity apart from occupation. *Brit J Prev Soc Med* 1967 ; 21 : 163-69.

ÉVALUATION DE L'ACTIVITÉ PHYSIQUE DES ENFANTS

- Adams J. Trends in physical activity and inactivity amongst US 14-18 year olds by gender, school grade and race, 1993-2003 : evidence from the youth risk behavior survey. *BMC Public Health* 2006 ; 6 : 57.
- Andersen LB, Haraldsdottir J. Tracking of cardiovascular disease risk factors including maximal oxygen uptake and physical activity from late teenager to adulthood : an 8-year follow-up study. *J Intern Med* 1993 ; 234, 309-15.
- Argiropoulou EC, Michalopoulou M, Aggeloussis N, Avgerinos A. Validity and reliability of physical activity measures in greek high school age children. *J Sports Sci Med* 2004 ; 3,147-59
- Atkin LM, Davies PS. Diet composition and body composition in preschool children. *Am J Clin Nutr* 2000 ; 72 : 15-21.
- Australian Bureau of Statistics. Children's participation in cultural and leisure activities. (Catalogue No. 4901.0). Canberra : ABS, 2004.
- Bailey R, Olson J, Pepper S, et al. The level and tempo of children's physical activities : an observational study. *Med Sci Sports Exerc* 1995 ; 27 : 1033-41.
- Bailey RC, Olson J, Pepper SL, et al. The level and tempo of children's physical activities : an observational study. *Med Sci Sports Exerc* 1995 ; 27 : 1033-41.
- Baranowski T, Thompson WO, DuRant RH, et al. Observations on physical activity in physical locations : age, gender, ethnicity, and month effects. *Res Q Exerc Sport* 1993 ; 64 : 127-33
- Beets MW, Patton MM, Edwards S. The Accuracy of Pedometer Steps and Time during Walking in Children. *Med Sci Sports Exerc* 2005 ; 3 : 513-20.
- Bender JM, Brownson RC, Elliott MB, Haire-Joshu DL. Children's physical activity : using accelerometers to validate a parent proxy record. *Med Sci Sports Exerc* 2005 ; 37 : 1049-13.
- Benham-Deal T. Preschool children's accumulated and sustained physical activity. *Percept Mot Skills* 2005 ; 100 : 443-50.
- Beunen GP, Lefevre J, Philippaerts RM, et al. Adolescent correlates of adult physical activity : a 26-year follow-up. *Med Sci Sports Exerc* 2004 ; 36 : 1930-6.
- Biddle S, Cavill N, Sallis J. *Young and active ? Young people and health-enhancing physical activity* – evidence and implications. London : Health Education Authority, 1998.
- Booth ML, Okely AD, Chey T, Bauman A. The reliability and validity of the physical activity questions in the WHO health behaviour in schoolchildren (HBSC) survey : a population study. *Br J Sports Med* 2001 ; 35 : 263-7.
- Burdette HL, Whitaker RC, Daniels SR. Parental report of outdoor playtime as a measure of physical activity in preschool-aged children. *Arch Pediatr Adolesc Med* 2004 ; 158 : 353-7.
- Caius N, Benefice E. Habitudes alimentaires, activité physique et surpoids chez des adolescents. *Rev Epidemiol Sante Publique* 2002 ; 50 : 531-42.

- Choquet M et al. *L'activité sportive à l'adolescence. Les troubles et conduites associées. Rapport remis au Ministère de la Jeunesse et des Sports.* Marly-le Roi : INJEP, 2000.
- de Vries SI, Bakker I, Hopman-Rock M, et al. Clinimetric review of motion sensors in children and adolescents. *J Clin Epidemiol* 2006 ; 59 : 670-80.
- Dowda M, Pate RR, Trost SG, et al. Influences of preschool policies and practices on children's physical activity. *J Community Health* 2004 ; 29 : 183-96.
- DuRant RH, Baranowski T, Davis H, et al. Reliability and variability of heart rate monitoring in 3-, 4-, or 5-yr old children. *Med Sci Sports Exerc* 1992 ; 24, 265-71.
- Durant RH, Baranowski T, Puhl J, et al. Evaluation of the Children's Activity Rating Scale (CARS) in young children. *Med Sci Sports Exerc* 1993a ; 25 : 1415-21.
- DuRant, RH, Baranowski T, Davis H, et al. Reliability and variability of indicators of heart rate monitoring in children. *Med Sci Sports Exerc* 1993c ; 25, 389-95.
- Eisenmann JC, Strath SJ, Shadrick D, et al. Validity of uniaxial accelerometry during activities of daily living in children. *Eur J Appl Physiol* 2004 ; 91 : 259-63.
- Ekelund U, Jan Åman J, Westerterp K. Is the ArteACC Index a Valid Indicator of Free-Living Physical Activity in Adolescents ? *Obes Res* 2003 ; 11 : 793-801.
- Ekelund U, Sjöström M, Yngve A, et al. Physical activity assessed by activity monitor and doubly labeled water in children. *Med Sci Sports Exerc*. 2001 ; 33 : 275-81.
- Fairweather SC, Reilly JJ, Grant S, et al. Using the Computer Science and Applications (CSA) activity monitor in children, *Pediatric Exerc Sci* 1999 ; 11, 413-20.
- Finn K, Johannsen N, Specker B. Factors associated with physical activity in preschool children. *J Pediatr* 2002 ; 140 : 81-5.
- Gavarry O, Falgairette G. Habitual physical activity during growth. *Can J Appl Physiol* 2004 ; 29 : 201-20.
- Going SL, Harrel J, Stewart D, et al. Physical activity assessment in American Indian schoolchildren in the Pathways study. *Am J Clin Nutr* 1999 ; 69 : 788S-95.
- Goran MI, Nagy TR, Gower BA, et al. Influence of sex, seasonality, ethnicity, and geographic location on the components of total energy expenditure in young children : implications for energy requirements. *Am J Clin Nutr* 1998 ; 68 : 675-82.
- Hallal PC, Wells JC, Reichert FF, et al. Early determinants of physical activity in adolescence : prospective birth cohort study. *Br Med J* 2006 ; 332 : 1002-7.
- Harro M. Validation of a questionnaire to assess physical activity of children ages 4-8 years. *Res Q Exerc Sport*, 1997 ; 4 : 259-68.
- Harvey-Berino J, Rourke J. Obesity prevention in preschool native-american children : a pilot study using home visiting. *Obes Res*. 2003 ; 11 : 606-11
- Hussey J, Gromley J, Bell C. Physical activity in Dublin schoolchildren aged 7-9 years. *Br J Sports Med* 2002 ; 36 : e2-003.
- Iannotti RJ, Claytor RP, Horn TS, Chen R. Heart Rate Monitoring as a Measure of Physical Activity in Children. *Med Sci Sports Exerc* 2004 ; 36 : 1964-71.
- Jackson DM, Reilly JJ, Kelly LA, et al. Objectively measured physical activity in a representative sample of 3- to 4-year-old children. *Obes Res* 2003 ; 11 : 420-5.
- Janz KF, Witt J, Mahoney LT. The stability of children's physical activity as measured by accelerometry and self-report. *Med Sci Sports Exerc* 1995 ; 27,1326-32.
- Janz KF. Validation of the CSA accelerometer for assessing children's physical activity. *Med Sci Sports Exerc* 1994 ; 26, 369-75.
- Klesges LM, Baranowski T, Beech B, et al. Social desirability bias in self-reported dietary, physical activity and weight concern measures in 8-to 10-year-old African-American girls : results from the Girls Health Enrichment Multisite Studies (GEMS). *Prev Med* 2004 ; 38 : S78-87.
- Kozuka N, Koo M, Allison KR, et al. The relationship between sedentary activities and physical inactivity among adolescents : results from the Canadian Community Health Survey. *J Adolesc Health* 2006 ; 39 : 515-22.
- Kohl HW III, Fulton JE, Caspersen CJ. Assessment of physical activity among children and adolescents : a review and synthesis. *Prev Med*. 2000 ; 31 : S54-76.
- Kraut A, Melamed S, Gofer D, Fromm P. Effect of School Age Sports on Leisure Time Physical Activity in Adults : The CORDIS Study. *Med Sci Sports Exerc* 2003 ; 12 : 2038-42.

- Lazzer S, Meyer F, Meyer M, et al. Évaluation de l'activité physique habituelle des adolescents en surpoids ou obèses. *Presse Med* 2004 ; 33 : 1255-9.
- Livingstone MB, Coward WA, Prentice AM, et al. Daily energy expenditure in free-living children : comparison of heart-rate monitoring with the doubly labeled water (2H218O) method. *Am J Clin Nutr* 1992 ; 56 : 343-52.
- Logan N, Reilley J, Grant S, Payton J. Resting heart rate definition and its effect on apparent levels of physical activity in young children. *Med Sci Sports Exerc* 2000 ; 32 : 162-6.
- Manios Y, Kafatos A, Markakis G. Physical activity of 6-year-old children : Validation of two proxy reports. *Pediatr Exerc Sci* 1998 ; 10 : 176-88.
- Manios Y. Design and descriptive results of the «Growth, Exercise and Nutrition Epidemiological Study In preSchoolers» : the GENESIS study. *BMC Public Health* 2006 ; 6 : 32.
- McManus A, Armstrong N. Patterns of physical activity among primary schoolchildren. In Ring F (Ed.) *Children in sport*, Bath University press, Bath, 1995 : 17-23
- Michaud PA, Caudey M, Narring F, Schutz Y. Assessment of physical activity with a pedometer and its relationship with VO2max among adolescents in Switzerland. *Soz Präventivmed* 2002 ; 47 : 107-15.
- Moore LL, Gao D, Bradlee ML, et al. Does early physical activity predict body fat change throughout childhood ? *Prev Med* 2003 ; 37 : 10-7.
- Moreau A, Pichault A, Truchot G. Le sport chez les jeunes de 12 à 17 ans. *Stat-Info Ministère des sport* 2002 ; 4 : 1-4. Les résultats complets sont disponibles sur le site Internet : www.jeunesse-sports.gouv.fr.
- Nilsson A, Ekelund U, Yngve A, Sjöström M. Levels of physical activity in children and adolescents in comparison with existing recommendations for health. *Med Sci Sports Exerc* 2002 ; 34 : S142.
- Nilsson A, Ekelund U, Yngve A, Sjöström M. Assessing physical activity in children with the CSA accelerometer : effect of different time sampling intervals and placements. *Pediatr Exerc Sci* 2002 ; 14 : 87-96.
- Noland M, Danner F, DeWalt K, et al. The measurement of physical activity in young children. *Res Q Exerc Sport* 1990 ; 61 : 146-53.
- Observatoire Régional de la Santé d'Alsace. *La santé des enfants de 6 ans en Alsace : analyse du questionnaire Nutrition Mode de vie – Visites médicales d'admission 2001/2002*. Strasbourg : ORSAL, avr 2004.
- O'Hara NM, Baranowski T, Simons-Morton BG, et al. Validity of the observation of children's physical activity. *Res Q Exerc Sport* 1989 ; 60 : 42-7.
- Pangrazi RP, Corbin CB, Welk GJ. Physical activity for children and youth. *J Phys Educ Rec Dance* 1996 ; 67 : 38-43.
- Parsons TJ, Power C, Manor O. Longitudinal physical activity and diet patterns in the 1958 British Birth Cohort. *Med Sci Sports Exerc* 2006 ; 38 : 547-54.
- Pate R. Critique of existing guidelines for physical activity in young people. In Biddle S, Sallis J, Cavill N (Eds) *Young and active ? Young people and health-enhancing physical activity – evidence and implications*. Health Education Authority, London, Great Britain, 1998 : 162-76.
- Pate RR, Baranowski T, Dowda M, Trost SG. Tracking of physical activity in young children. *Med Sci Sports Exerc*. 1996 ; 28 : 92-6.
- Pate RR, Freedson PS, Sallis JF, et al. Compliance with Physical Activity Guidelines : Prevalence in a Population of Children and Youth. *Ann Epidemiol* 2002 ; 12 : 303-8.
- Pate RR, Pfeiffer KA, Trost SG, et al. Physical activity among children attending preschools. *Pediatrics* 2004 ; 114 : 1258-63.
- Prochaska JJ, Sallis JF, Long B. A physical activity screening measure for use with adolescents in primary care. *Arch Pediatr Adolesc Med* 2001 ; 155 : 554-9.
- Puyau MR, Adolph AL, Firoz AV, Butte NF. Validation and calibration of physical activity monitors in children. *Obe Res*. 2002 ; 10 : 150-7.
- Puyau MR, Adolph AL, Vohra FA, et al. Prediction of Activity Energy Expenditure Using Accelerometers in Children. *Med Sci Sports Exerc* 2004 ; 36 : 1625-31.
- Ramirez-Marrero F, Smith BE, Kirby TE, et al. Evaluation of a step-counter during treadmill walking in 7-12 year old African-American children. *J Natl Black Nurses Assoc* 2002 ; 13 : 1-5.
- Reilly JJ, Coyle J, Kelly L, et al. An objective method for measurement of sedentary behavior in 3- to 4 year olds. *Obes Res* 2003 ; 11 : 1155-8.
- Reilly JJ, Jackson DM, Montgomery C, et al. Total energy expenditure and physical activity in young Scottish children : mixed longitudinal study. *Lancet* 2004 ; 363 : 211-2.

Riddoch C, Boreham C. The health-related physical activity of children. *Sports Med* 1995 ; 19 : 86-102.

Riddoch C, Boreman C. The health-related physical activity of children. *Res Q Exerc Sport* 1995a ; 65 : 84-8.

Ridgers ND, Stratton G, Fairclough SJ. Assessing physical activity during recess using accelerometry. *Prev Med* 2005 ; 41 : 102-7.

Ridley K, Olds T, Hill A. A multimedia activity recall for children and adolescents (MARCA) — description and validation. *Australian conference of science and medicine in sport 2002 : Sports medicine and science at the extremes*. 12-16 October 2002, Melbourne.

Rodriguez G, Beghin L, Michaud L, et al. Comparison of the TriTrac-R3D accelerometer and a self-report activity diary with heart-rate monitoring for the assessment of energy expenditure in children. *Br J Nutr* 2002 ; 87 : 623-31.

Rowe PJ, Schuldheisz JM, van der Mars H. Validation of SOFIT for measuring physical activity of first- to eighth-grade students. *Pediatr Exerc Sci* 1997 ; 9 : 136-49.

Rowlands AV, Thomas PW, Eston RG, Topping R. Validation of the RT3 triaxial accelerometer for the assessment of physical activity. *Med Sci Sports Exerc* 2004 ; 36 : 518-24

Salbe AD, Fontvieille AM, Harper IT, Ravussin E. Low levels of physical activity in 5-year-old children. *J Pediatr* 1997 ; 131 : 423-9.

Sallis JF, Berry CC, Broyles SL, McKenzie TL, Nader PR. Variability and tracking of physical activity over two years in young children. *Med Sci Sports Exerc* 1995 ; 27 : 1042-9.

Sallis JF, Buono MJ, Roby JJ, et al. Seven-day recall and other physical activity self-reports in children and adolescents. *Med Sci Sports Exerc* 1993 ; 25 : 99-108.

Sallis JF, Buono MJ, Roby JJ, et al. The Caltrac accelerometer as a physical activity monitor for school-age children. *Med Sci Sports Exerc* 1990 ; 22 : 698-703.

Sallis JF, Strikmiller PK, Harsha DW, et al. Validation of interviewer- and self-administered physical activity checklists for fifth grade students. *Med Sci Sports Exerc* 1996 ; 28 : 840-51.

Sallis JF. Self-report measures of children's physical activity. *J School Health* 1991 ; 61,215-9.

Sallo M, Silla R. Physical activity with moderate to vigorous intensity in pre -school and first grade children. *Ped Exerc Sci*. 1997 ; 9 : 44-54

Schmitz KH, et al. Reliability and validity of a brief questionnaire to assess television viewing and computer use by middle school children. *J Sch Health* 2004 ; 74 : 370-7.

Simon C, Klein C, Wagner A. La sédentarité des enfants et des adolescents, un enjeu de santé publique. *J Pédiatrie Puériculture* 2005 ; 18 : 217-23.

Simons-Morton BG, Taylor WC, Huang IW. Validity of the physical activity interview and Caltrac with preadolescent children. *Res Q Exerc Sport* 1994 ; 65 : 84-8.

Sirard JR, Pate RR. Physical activity assessment in children and adolescents. *Sports Med* 2001 ; 31 : 439-54.

Spadano JL, Bandini LG, Must A, et al. Longitudinal changes in energy expenditure in girls from late childhood through midadolescence. *Am J Clin Nutr* 2005 ; 5 : 1102-9.

Tammelin T, Näyhä S, Laitinen J, et al. Physical activity and social status in adolescence as predictors of physical inactivity in adulthood. *Prev Med* 2003 ; 37 : 375-81.

Telama R, Yang X, Viikari J, et al. Physical activity from childhood to adulthood : a 21-year tracking study. *Am J Prev Med* 2005 ; 28 : 267-73.

Tennefors C, Coward WA, Hernell O, et al. Total energy expenditure and physical activity level in healthy young Swedish children 9 or 14 months of age. *Eur J Clin Nutr* 2003 ; 57 : 647-53.

Tomkinson GR, Leger LA, Olds TS, Cazorla G. Secular trends in the performance of children and adolescents (1980-2000) : an analysis of 55 studies of the 20m shuttle run test in 11 countries. *Sports Med* 2003 ; 33 : 285-300.

Tremblay MS, Inman JW, Willms JD. Preliminary evaluation of a video questionnaire to assess activity levels of children. *Med Sci Sports Exerc* 2001 ; 33 : 2139-44.

Treuth MS, Hou N, Young DR, Maynard LM. Validity and reliability of the Fels physical activity questionnaire for children. *Med Sci Sports Exerc* 2005 ; 37 : 488-95.

Treuth MS, Sherwood NE, Butte NF, et al. Validity and reliability of activity measures in African-American girls for GEMS. *Med Sci Sports Exerc* 2003 ; 35 : 532-9.

Trost SG, Pate RR, Freedson PS, et al. Using objective physical activity measure with youth : How many days of monitoring are needed. *Med Sci Sports Exerc* 2000 ; 35, 426-31.

- Trudeau F, Laurencelle L, Shephard RJ. Tracking of physical activity from childhood to adulthood. *Med Sci Sports Exerc* 2004 ; 36 : 1937-43.
- Tudor-Locke C, Pangrazi RP, Corbin CB, et al. BMI-referenced standards for recommended pedometer-determined steps/day in children. *Prev Med* 2004 ; 38 : 857-64.
- Van Covering P, Harnack L, Schmitz K, et al. Feasibility of Using Accelerometers to Measure Physical Activity in Young Adolescents. *Med Sci Sports Exerc* 2005 ; 37 : 867-71.
- Welk GJ, Corbin CB. The validity of the Tritrac R-3D activity monitor for assessment of physical activity. *Res Q Exerc Sport* 1995 ; 66 : 202-9.
- Welk GJ, Dziewaltowski DA, Hill JL. Comparison of the computerized ACTIVITYGRAM instrument and the previous day physical activity recall for assessing physical activity in children. *Res Q Exerc Sport* 2004 ; 75 : 370-80.
- Weston AT, Petosa R, Pate RR. Validation of an instrument for measurement of physical activity in youth. *Med Sci Sports Exerc* 1997 ; 29,138-43.
- Williams CL, Carter BJ, Wynder EL. Prevalence of selected cardiovascular and cancer risk factors in a pediatric population : the "Know Your Body" Project, New York. *Prev Med* 1981 ; 10 : 235-50.
- Zask A, van Beurden E, Barnett L, et al. Active School Playgrounds-Myth or Reality. Results of the « Move it Groove it » Project. *Prev Med* 2001 ; 33 : 402-8.

ÉVALUATION DE L'ACTIVITÉ PHYSIQUE DES SENIORS

- Black AE, Coward WA, Cole TJ, Prentice AM. Human energy expenditure in affluent societies : an analysis of 574 doubly-labelled water measurements. *Eur J Clin Nutr* 1996 ; 50 : 72-92.
- Bonnefoy M, Kostka T, Berthouze SE, Lacour JR. Validation of a physical activity questionnaire in the elderly. *Eur J Appl Physiol Occup Physiol* 1996 ; 74 : 528-33.
- Bonnefoy M, Normand S, Pachiardi C, et al. Simultaneous validation of ten physical activity questionnaires in older men : a doubly labeled water study. *J Am Geriatr Soc* 2001 ; 49 : 28-35.
- Brach JS, FitzGerald S, Newman AB, et al. Physical activity and functional status in community-dwelling older women. *Arch Intern Med* 2003 ; 163 : 2565-71.
- Chan M, Campo E, Laval E, Estève D. Validation of a remote monitoring system for the elderly : application to mobility measurements. *Technol Health Care* 2002 ; 10 : 391-9.
- De Abajo S, Larriba R, Marquez S. Validity and reliability of the Yale Physical Activity Survey in Spanish elderly. *J Sports Med Phys Fitness* 2001 ; 41 : 479-85
- Dinger MK, Oman RF, Taylor EL, Vesely SK, Able J. Stability and convergent validity of the Physical Activity Scale for the Elderly (PASE). *J Sports Med Phys Fitness* 2004 ; 44 : 186-92.
- Dipietro L, Caspersen CJ, Ostfeld AM, Nadel ER. A survey for assessing physical activity among older adults. *Med Sci Sports Exerc* 1993 ; 25 : 628-42.
- Dressen C, Perrin H. La santé des personnes âgées de 60 à 75 ans. In : Guilbert P, Baudier F, Gautier A (dir). *Baromètre Santé 2000*. Vol 2 : Résultats. Vanves : CFES, 2001.
- Dubbert PM, Vander Weg MW, Kirchner KA, Shaw B. Evaluation of the 7-Day Physical Activity Recall in Urban and Rural Men. *Med Sci Sports Exerc* 2004 ; 36 : 1646-54.
- Dvorak RV, Tchernof A, Starling RD, et al. Respiratory fitness, free living physical activity, and cardiovascular disease risk in older individuals : a doubly labeled water study. *J Endocrinol Metab* 2000 ; 85 : 957-63.
- Glass TA, de Leon CM, Marottoli RA, Berckman LF. Population based study of social and productive activities as predictors of survival among elderly Americans. *Br J Med* 1999 ; 319 : 478-83.
- Hansen RD, Allen BJ. Habitual physical activity, anabolic hormones, and potassium content of fat-free mass in postmenopausal women. *Am J Clin Nutr* 2002 ; 75 : 314-20.
- Jouandou PO, Constans T. Pourquoi les sujets âgés sportifs arrêtent-ils le sport ? *L'année gériatrique* 2005 ; 19 : 61-70.
- Kochersberger G, McConnell E, Kuchibhatla MN, Pieper C. The reliability, validity, and stability of a measure of physical activity in the elderly. *Arch Phys Med Rehabil*.1996 ; 77 : 793-5

La pratique des activités physiques et sportive en France. Résultats de l'enquête menée en 2003 dans le cadre d'un partenariat entre le ministère de la Jeunesse, des Sports et de la Vie Associative, le ministère de la Culture et de la Communication et l'INSEE. Paris : MJSVA & INSEP, 2005.

Lawlor DA, Taylor M, Bedford C, Ebrahim S. Is housework good for health ? Levels of physical activity and factors associated with activity in elderly women. *J Epidemiol Community Health* 2002 ; 56 : 473-8.

Le Masurier GC, Tudor-Locke C. Comparison of pedometer and accelerometer accuracy under controlled conditions. *Med Sci Sports Exerc* 2003 ; 35 : 867-71.

Lyons GM, Culhane KM, Hilton D, et al. description of an accelerometer-based mobility monitoring technique. *Med Eng Phys* 2005 ; 27 : 497-504.

Melanson EL, Knoll JR, Bell ML, et al. Commercially available pedometers : considerations for accurate step counting. *Prev Med* 2004 ; 39 : 361-8.

OMS. *Viellir en restant actif. Cadre d'orientation.* WHO/NMH/NPH/02.8. Genève : OMS, 2002.

Pols MA, Peeters PH, Kemper HC, Collette HJ. Repeatability and relative validity of two physical activity questionnaires in elderly women. *Med Sci Sports Exerc* 1996 ; 28 : 1020-5.

Pols MA, Peeters PH, Ocke MC, et al. Estimation of reproducibility and relative validity of the questions included in the EPIC Physical Activity Questionnaire. *Int J Epidemiol* 1997 ; 26 : S181-9.

Scanail C, Carew S, Barralon P, et al. Review of approaches to mobility telemonitoring of the elderly in their living environment. *Ann Biomed Engineering* 2006 ; 17 : 547-63.

Scanail C, Lyons GM. Long-term telemonitoring of mobility trends of elderly people using SMS messaging. *IEEE Trans Inf Technol Biomed* 2006 ; 10 : 412-3

Schoeller DA, van Santen E. Measurement of energy expenditure in humans by doubly labeled water method. *J Appl Physiol* 1982 ; 53 : 955-9.

Schuit AJ, Schouten EG, Westerterp KR, Saris WH. Validity of the Physical Activity Scale for the Elderly (PASE) : according to energy expenditure assessed by the doubly labeled water method. *J Clin Epidemiol* 1997 ; 50 : 541-6.

Schuler PB, Richardson MT, Ochoa P, Wang MQ. Accuracy and repeatability of the Yale physical activity survey in assessing physical activity of older adults. *Percept Mot Skills* 2001 ; 93 : 163-77.

Seale JL, Klein G, Friedmann J, et al. Energy expenditure measured by doubly labeled water, activity recall, and diet records in the rural elderly. *Nutrition* 2002 ; 18 : 568-73.

Starling RD, Matthews DE, Ades PA, Poehlman ET. Assessment of physical activity in older individuals : a doubly labeled water study. *J Appl Physiol* 1999 ; 86 : 2090-6.

Stel VS, Smit JH, Pluijm SM, et al. Comparison of the LASA Physical Activity Questionnaire with a 7-day diary and pedometer. *J Clin Epidemiol* 2004 ; 57 : 252 -8.

Struck BD, Ross KM. Health promotion in older adults. Prescribing exercise for the frail and home bound. *Geriatrics* 2006 ; 61 : 22-7.

Taylor-Piliae RE, Norton LC, Haskell WL, et al. Validation of a New Brief Physical Activity Survey among Men and Women Aged 60-69 Years. *Am J Epidemiol* 2006 ; 164 : Jul 13.

The importance of physical activity for older adults. Michigan Governor's Council on Physical Fitness, health and Sports. Michigan Department of Community Health, July, 2002.

Washburn RA, Ficker JL. Physical Activity Scale for the Elderly (PASE) : the relationship with activity measured by a portable accelerometer. *J Sports Med Phys Fitness* 1999 ; 39 : 336-40.

Washburn RA, Smith KW, Jette AM, Janney CA. The Physical Activity Scale for the Elderly (PASE) : development and evaluation. *J Clin Epidemiol* 1993 ; 46 : 153-62.

Westerterp KR. Daily physical activity and ageing. *Curr Opin Clin Nutr Metab Care* 2000 ; 3 : 485-8.

WHO. *Guidelines series for healthy ageing.1. The Heidelberg guidelines for promoting physical activity among older persons.* Ageing and Health Program, Division of Health Promotion, Education and Communication, 1996.

- Alberti KGGM, Zimmet PZ, for the WHO Consultation. Definition, diagnosis and classification of diabetes mellitus and its complications. Part I, Diagnosis and classification of diabetes mellitus. Provisional report of a WHO consultation. *Diabet Med* 1998 ;15 : 539-53.
- American Dietetic Association (ADA). Position of the American Dietetic Association : individual-, family-, school-, and community-based interventions for pediatric overweight. *J Am Diet Assoc* 2006 ; 106 : 925-45.
- Araiza P, Hewes H, Gashetewa C, et al. Efficacy of a pedometer-based physical activity program on parameters of diabetes control in type 2 diabetes mellitus. *Metabolism* 2006 ; 55 : 1382-7.
- Balkau B, Vernay M, Mhamdi L, et al. The incidence and persistence of the NCEP (National Cholesterol Education Program) metabolic syndrome. The French D.E.S.I.R. study. *Diabetes Metab* 2003 ; 29 : 526-32.
- Bouchard C, Després JP, Tremblay A. Exercise and obesity. *Obes Res* 1993 ; 1 : 133-47.
- Després JP, Lamarche B. Effects of diet and physical activity on adiposity and body fat distribution : implications for the prevention of cardiovascular disease. *Nutr Res Rev* 1993 ; 6 : 137-59.
- Doak CM, Visscher TL, Renders CM, Seidell JC. The prevention of overweight and obesity in children and adolescents : a review of interventions and programmes. *Obes Rev* 2006 ; 7 : 111-3.
- Donnelly JE, Smith B, Jacobsen DJ, et al. The role of exercise for weight loss and maintenance. *Best Pract Res Clin Gastroenterol* 2004 ; 18 : 1009-29.
- Executive summary of the third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel III). *J Am Med Assoc* 2001 ; 285 : 2846-497.
- Garrow JS, Summerbell CD. Meta-analysis : effect of exercise, with or without dieting, on the body composition of overweight subjects. *Eur J Clin Nutr* 1995 ; 49 : 1-10
- Garrow JS. Exercise in the treatment of obesity : a marginal contribution. *Int J Obes Rel Metab Dis* 1995 ; 19 : S216-9.
- Gomila S, Dallongeville J. Epidémiologie du syndrome métabolique en France. *Med Nutr* 2003 ; 39 : 89-94.
- Halbert JA, Silagy CA, Finucane P, et al. Exercise training and blood lipids in hyperlipemic and normolipemic adults : a meta-analysis of randomised, controlled trials. *Eur J Clin Nutr* 1999 ; 53 : 514-22.
- Hambrecht R, Niebauer J, Marburger C, et al. Various intensities of leisure-time physical activity in patients with coronary artery disease : effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. *J Am Coll Cardiol* 1993 ; 22 : 468-7.
- Hardman AE, Herd S.L. Exercise and postprandial lipid metabolism. *Proc Nutr Soc* 1998 ; 57 : 63-72.
- Hayashi T, Wojtaszewski JPF, Goodyear LJ. Exercise regulation of glucose transport in skeletal muscle. *Am J Physiol* 1997 ; 273 : E1039-E51.
- Herbst A, Bachran R, Kapellen T, Holl RW. Effects of regular physical activity on control of glycemia in pediatric patients with type 1 diabetes mellitus. *Arch Pediatr Adolesc Med* 2006 ; 160 : 573-7.
- Jeffery RW, Wing RR, Sherwood NE, Tate DF. Physical activity and weight loss : does prescribing higher physical activities goals improve outcome ? *Am J Clin Nutr* 2003 ; 78 : 684-9.
- Katzmarzyk PT, Leon AS, Wilmore JH, et al. Targeting the metabolic syndrome with exercise : evidence from the HERITAGE Family Study. *Med Sci Sports Exerc* 2003 ; 35 : 1703-9.
- Klem ML, Wing RR, McGuire MT, et al. A descriptive study of individuals successful at long-term maintenance of substantial weight loss. *Am J Clin Nutr* 1997 ; 66 : 239-46.
- Lamarche B, Després JP, Pouliot MC, et al. Is body fat loss a determinant factor in the improvement of carbohydrate and lipid metabolism following aerobic exercise training in obese women ? *Metabolism* 1992 ; 41 : 1249-56.
- Ligtenberg PC, Blans M, Hoekstra JBL, et al. No effect of long-term physical activity on the glycemic control in type 1 diabetes patients : a cross-sectional study. *Neth J Med* 1999 ; 55 : 59-63.

- Miller WC, Koceja DM, Hamilton EJ. A meta-analysis of the past 25 years of weight loss research using diet, exercise or diet plus exercise intervention. *Int J Obes Relat Metab Disord* 1997 ; 21 : 941-7.
- Obesity : Preventing and Managing the Global Epidemic*. Report of a WHO Consultation on Obesity, Geneva, 3-5 June 1997. Geneva : World Health Organization ; 1997.
- Oppert JM, Rolland-Cachera MF. Prevalence, évolution dans le temps et conséquences économiques de l'obésité. *Médecine/sciences* 1998 ; 11 : 939-43.
- Plourde G. Preventing and managing pediatric obesity. Recommendations for family physicians. *Can Fam Physician* 2006 ; 52 : 322-8.
- Raynor DA, Phelan S, Hill JO, Wing RR. Television viewing and long-term weight maintenance : results from the National Weight Control Registry. *Obesity* 2006 ; 14 : 1816-24.
- Richardson CR, Brown BB, Foley S, et al. Feasibility of Adding Enhanced Pedometer Feedback to Nutritional Counseling for Weight Loss. *J Med Internet Res* 2005 ; 7 (5) : e56.
- Riddell MC, Perkins BA. Type 1 diabetes and vigorous exercise : applications of exercise physiology to patient management. *Can J Diabetes* 2006 ; 30 : 63-71.
- Riebe D, Blissmer B, Greene G, et al. Long-term maintenance of exercise and healthy eating behaviors in overweight adults. *Prev Med* 2005 ; 40 : 769-78.
- Ross R, Dagnone D, Jones PJ, et al. Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. *Ann Intern Med* 2000 ; 133 : 92-103.
- Ross R, Janssen I. Physical activity, total and regional obesity : dose-response considerations. *Med Sci Sports Exerc* 2001 ; 33 : S521-9.
- Saris WH, Blair SN, van Baak MA, et al. How much physical activity is enough to prevent unhealthy weight gain ? Outcome of the IASO 1st Stock Conference and consensus statement. *Obes Rev* 2003 ; 4 : 101-14
- Schmidt WD, Biber CJ, Kalscheuer LK. Effects of long versus short bout exercise on fitness and weight loss in overweight females. *J Am Coll Nutr* 2001 ; 20 : 494-501.
- Schneider PL, Bassett DR Jr, Thompson DL, et al. Effects of a 10,000 steps per day goal in overweight adults. *Am J Health Promot* 2006 ; 21 : 85-9.
- Slentz CA, Duscha BD, Johnson JL, et al. Effects of the amount of exercise on body weight, body composition, and measures of central obesity : STRRIDE—a randomized controlled study. *Arch Intern Med* 2004 ; 164 : 31-9.
- Stewart KJ, Bacher AC, Turner K, et al. Exercise and risk factors associated with metabolic syndrome in older adults. *Am J Prev Med* 2005 ; 28 : 9-18.
- The European Group for the Study of Insulin Resistance (EGIR). Frequency of the WHO metabolic syndrome in European cohorts, and an alternative definition of an Insulin resistance syndrome. *Diabetes Metab* 2002 ; 28 : 364-76.
- Thomas D, Elliott E, Naughton G. Exercise for type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2006 Jul 19 ; 3 : CD002968.
- Tremblay A, Fontaine E. Contribution of postexercise increment in glucose storage to variations in glucose-induced thermogenesis in endurance athletes. *Can J Physiol Pharmacol* 1985 ; 63 : 1165-9.
- Vischer UM. La prévention du diabète de type 2 : activité physique ou médicaments ? *Med Hyg* 2003 ; 61 : 1192-7.
- Waden J, Tikkanen H, Forsblom C, et al. Leisure time physical activity is associated with poor glycemic control in type 1 diabetic women : the FinnDiane study. *Diabetes Care* 2005 ; 28 : 777-82.
- Wahrenberg H, Bolinder J, Arner P. Adrenergic regulation of lipolysis in human fat cells during exercise. *Eur J Clin Invest* 1991 ; 21 : 534-41.
- Watkins LL, Sherwood A, Feinglos M, et al. Effects of exercise and weight loss on cardiac risk factors associated with syndrome X. *Arch Intern Med* 2003 ; 163 : 1889-95.
- Yusuf S, Hawken S, Ounpuu S, et al. Obesity and the risk of myocardial infarction in 27 000 participants from 52 countries : a case-control study. *Lancet* 2005 ; 366 : 1640-49
- Ziegler O, Quilliot T. Prise en charge de l'obésité de l'adulte. *Rev Prat* 2005 ; 55 : 1437-52

- Adams J, Ogola G, Stafford P, et al. High-intensity interval training for intermittent claudication in a vascular rehabilitation program. *J Vasc Nurs* 2006 ; 24 : 46-9.
- Ades PA, Savage PD, Tischler MD, et al. Determinants of disability in older coronary patients. *Am Heart J* 2002 ; 143 : 151-6.
- Belardinelli R, Georgiou D, Cianci G, Purcaro A. Randomized, controlled trial of long term moderate exercise training in chronic heart failure : effects on functional capacity, quality of life, and clinical outcome. *Circulation* 1999 ; 99 : 1173-82.
- Bendermacher BL, Willigendael EM, Teijink JA, Prins MH. Supervised exercise therapy versus non-supervised exercise therapy for intermittent claudication. *Cochrane Database Syst Rev* 2006 Apr 19 ; (2) : CD005263.
- Benoit E, Berzin A, Foratier V, et al. Prise en charge des facteurs de risque cardiovasculaire après un infarctus du myocarde. *Rev Med Assur Maladie* 2005 ; 36 : 207-16.
- Briffa TG, Eckermann SD, Griffiths AD, et al. Cost-effectiveness of rehabilitation after an acute coronary event : a randomised controlled trial. *Med J Aust* 2005 ; 183 : 450-5
- Briffa TG, Maiorana A, Sheerin NJ, et al. Physical activity for people with cardiovascular disease : recommendations of the National Heart Foundation of Australia. *Med J Aust* 2006 ; 184 : 71-5.
- Brochu M, Poehlman ET, Savage P, et al. Modest effects of exercise training alone on coronary risk factors and body composition in coronary patients. *J Cardiopulm Rehabil* 2000 ; 20 : 180-8.
- Bulmer AC, Coombes JS. Optimising exercise training in peripheral arterial disease. *Sports Med* 2004 ; 34 : 983-1003.
- Clark AM, Hartling L, Vandermeer B, McAlister FA. Meta-analysis : secondary prevention programs for patients with coronary artery disease. *Ann Intern Med* 2005 ; 143 : 659-72.
- Collins E, Langbein WE, Dilan-Koetje J, et al. Effects of exercise training on aerobic capacity and quality of life in individuals with heart failure. *Heart Lung* 2004 ; 33 : 154-61.
- Corvera-Tindel T, Doering LV, et al. Effects of a home walking exercise program on functional status and symptoms in heart failure. *Am Heart J* 2004 ; 147 : 339-46.
- Dafoe W, Huston P. Current trends in cardiac rehabilitation. *Can Med Assoc J* 1997 ; 156 : 527-32.
- Delagardelle C, Feiereisen P. Strength training for patients with chronic heart failure. *Eura Medicophys* 2005 ; 41 : 57-65.
- Douard H, Vuillemin C, Bordier P, et al. Profils tensionnels normaux à l'effort selon l'âge, le sexe et les protocoles. *Arch Mal Coeur* 1994 ; 87 : 311-318
- Elley R, Bagrie E, Arroll B. Do snacks of exercise lower blood pressure ? A randomised crossover trial. *N Z Med J* 2006 ; 119 : U1996.
- Fagard RH. Exercise characteristics and the blood pressure response to dynamic physical training. *Med Sci Sports Exerc* 2001 ; 6 : S484-92.
- Fagard RH. Physical activity in the prevention and treatment of hypertension in the obese. *Med Sci Sports Exerc* 1999 ; 31 : S624-30.
- Franklin BA, Bonzheim K, Gordon S, Timmis GC. Safety of medically supervised outpatient cardiac rehabilitation exercise therapy : a 16-year follow-up. *Chest* 1998 ; 114 : 902-6.
- Gardner AW, Katzel LI, Sorkin JD, et al. Exercise rehabilitation improves functional outcomes and peripheral circulation in patients with intermittent claudication : a randomized controlled trial. *J Am Geriatr Soc* 2001 ; 49 : 755-62
- Gardner AW, Montgomery PS, Flinn WR, Katzel LI. The effect of exercise intensity on the response to exercise rehabilitation in patients with intermittent claudication. *J Vasc Surg* 2005 ; 42 : 702-9.
- Gardner AW, Poehlman ET. Exercise rehabilitation programs for the treatment of claudication pain. A meta-analysis. *JAMA* 1995 ; 274 : 975-80.
- Giannuzzi P, Temporelli PL, Corra U, et al. Antiremodeling effect of long-term exercise training in patients with stable chronic heart failure : results of the Exercise in Left Ventricular Dysfunction and Chronic Heart Failure (ELVD-CHF) Trial. *Circulation* 2003 ; 108 : 554-9.
- Hagberg JM, Montain SJ, Martin WH, Ehsani AA. Effect of exercise training in 60- to 69-year-old persons with essential hypertension. *Am J Cardiol* 1989 ; 64 : 348-53.
- Hagberg JM, Park JJ, Brown MD. The role of exercise training in the treatment of hypertension : an update. *Sports Med* 2000 ; 30 : 193-2006.

Hambrecht R, Fiehn E, Weigl C, et al. Regular physical exercise corrects endothelial dysfunction and improves exercise capacity in patients with chronic heart failure. *Circulation* 1998 ; 98 : 2709-15.

Hambrecht R, Walther C, Möbius-Winkler S, et al. Percutaneous coronary angioplasty compared with exercise training in patients with stable coronary artery disease. A randomized trial. *Circulation* 2004 ; 109 : 1371-8.

Hung C, Daub B, Black B, et al. Exercise training improves overall physical fitness and quality of life in older women with coronary artery disease. *Chest* 2004 ; 126 : 1026-31.

Jancik J, Siegelova J, Homolka P, et al. Exercise training in patients with chronic coronary heart disease. *Scripta Medica* 2002 ; 75 : 151-6.

Kavanagh T, Mertens DJ, Hamm LF, et al. Peak oxygen intake and cardiac mortality in women referred for cardiac rehabilitation. *J Am Coll Cardiol* 2003 ; 42 : 2139-4.

Kavanagh T, Mertens DJ, Hamm LF, et al. Prediction of long-term prognosis in 12 169 men referred for cardiac rehabilitation. *Circulation* 2002 ; 106 : 666 -7

Keller G, Tran ZV. Aerobic exercise and normotensive adults : a meta-analysis. *Med Sci Sport Exerc.* 1995 ; 27 : 1371-7

Kelley G, McClellan P. Antihypertensive effects of aerobic exercise : a brief meta-analytic review of randomized controlled trials. *Am J Hypertens* 1994 ; 7 : 115-9.

Kiilavuori K, Naveri H, Salmi T, Harkonen M. The effect of physical training on skeletal muscle in patients with chronic heart failure. *Eur J Heart Fail* 2000 ; 2 : 53-63.

Kokkinos P, Narayan P, Collier JA, et al. Effects of regular exercise on blood pressure and left ventricular hypertrophy in africanamerican men with severe hypertension. *N Engl J Med* 1995 ; 333 : 1462-7.

Kokkinos P, Pittaras A, Manolis A, et al. Exercise capacity and 24-h blood pressure in prehypertensive men and women. *Am J Hypertens* 2006 ; 19 : 251-8.

Leng GC, Fowler B, Ernst E. Exercise for intermittent claudication. *Cochrane Database Syst Rev* 2000 ; (2) : CD000990.

Leon AS, Franklin BA, Costa F, et al. Cardiac rehabilitation and secondary prevention of coronary heart disease. *Circulation* 2005 ; 111 : 369-76.

Levinger I, Bronks R, Cody DV, et al. The effect of resistance training on left ventricular function and structure of patients with chronic heart failure. *Int J Cardiol* 2005 ; 105 : 159-63.

Lipchenko AA, Fomin IL. Use of physical training in the rehabilitation of patients with post-infarct heart failure. *Kardiologija* 1987, 27 : 48-52.

Matsusaki M, Ikeda M, Tashiro E, et al. Influence of work load on the antihypertensive effect of exercise. *Clin Exp Pharmacol Physiol* 1992 ; 19 : 471-9.

McDermott MM, Liu K, Ferrucci L, et al. Physical performance in peripheral arterial disease : a slower rate of decline in patients who walk more. *Ann Intern Med* 2006 ; 144 : 10-20.

McKelvie RS, Teo KK, Roberts R, et al. Effects of exercise training in patients with heart failure : the Exercise Rehabilitation Trial (EXERT). *Am Heart J* 2002 ; 144 : 23-30.

Moreira WD, Fuchs FD, Ribeiro JP, Appel LJ. The effects of two aerobic training intensities on ambulatory blood pressure in hypertensive patients : results of a randomized trial. *J Clin Epidemiol* 1999 ; 52 : 637-42.

Mundal R, Kjeldsen SE, Sandvik L, et al. Exercise blood pressure predicts mortality from myocardial infarction. *Hypertension* 1996 ; 27 : 324-329

Nelson L, Esler MD, Jennings GL, Korner PI. Effect of changing levels of physical activity on blood pressure and haemodynamics in essential hypertension. *Lancet* 1986 ; 2 : 473-6.

Ng PW, Hollingsworth SJ, Luery H, et al. Intermittent claudication : exercise-increased walking distance is not related to improved cardiopulmonary fitness. *Eur J Vasc Endovasc Surg* 2005 ; 30 : 391-4.

Pescatello LS, Franklin BA, Fagard R, et al. American College of Sports Medicine position stand. Exercise and hypertension. *Med Sci Sports Exerc.* 2004 ; 3 : 533-553.

Piepoli MF, Davos C, Francis DP, et al. Exercise training meta-analysis of trials in patients with chronic heart failure (ExTraMATCH). *BMJ* 2004 ; 328 (7433) : 189.

Rogers MW, Probst MM, Gruber JJ, et al. Differential effects of exercise on blood pressure and cardiovascular responses to stress in borderline hypertensive humans. *J Hypertens* 1996 ; 14 : 1369-75.

- Rognmo O, Hetland E, Helgerud J, et al. High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease. *Eur J Cardiovasc Prev Rehabil* 2004 ; 11 : 216-2.
- Slordahl SA, Wang E, Hoff J, et al. Effective training for patients with intermittent claudication. *Scand Cardiovasc J* 2005 ; 39 : 244-9.
- Spronk S, Bosch JL, Veen HF, et al. Intermittent claudication : functional capacity and quality of life after exercise training or percutaneous transluminal angioplasty--systematic review. *Radiology* 2005 ; 235 : 833-42.
- Stahle A, Mattsson E, Ryden L, et al. Improved physical fitness and quality of life following training of elderly patients after acute coronary events. A 1 year follow-up randomized controlled study. *Eur Heart J* 1999 ; 20 : 1475-84.
- Stewart KJ, Hiatt WR, Regensteiner JG, Hirsch AT. Exercise training for claudication ; a review. *N Engl J Med* 2002 ; 347 : 1941-51.
- Stewart KJ, Turner KL, Bacher AC, et al. Are fitness, activity, and fatness associated with health- related quality of life and mood in older persons ? *J Cardiopulm Rehabil* 2003 ; 23 : 115-21.
- Takeda M, Tanaka K, Unno H, et al. Changes in aerobic capacity and coronary risk factors during long-term exercise training in women with ischemic heart disease : a 36-month follow-up. *Appl Human Sci* 1996 ; 15 : 115-21.
- Tashiro E, Miura S, Koga M et al. Crossover comparison between the depressor effects of low and high work rate exercise in mid hypertension. *Clin Exp Pharmacol Physiol* 1993 ; 20 : 689-96.
- Taylor RS, Brown A, Ebrahim S, et al. Exercise-based rehabilitation for patients with coronary heart disease : systematic review and meta-analysis of randomized controlled trials. *Am J Med* 2004 ; 116 : 682-92.
- Taylor-Tolbert N, Dengel D, Brown MD, Other AN. Ambulatory blood pressure after acute exercise in older men with essential hypertension. *Am J Hypertension* 2000 ; 13 : 44-51.
- Thompson PD, Buchner D, Pina IL, et al. Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease. *Circulation* 2003 ; 107 : 3109-16.
- Tokmakidis SP, Volaklis KA. Training and detraining effects of a combined-strength and aerobic exercise program on blood lipids in patients with coronary artery disease. *J Cardiopulm Rehabil* 2003 ; 23 : 193-200.
- Ueshima K, Suzuki T, Nasu M, et al. Effects of exercise training on left ventricular function evaluated by the Tei index in patients with myocardial infarction. *Circ J* 2005 ; 69 : 564-6.
- Whelton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure : A meta-analysis of randomized controlled trials. *Ann Int Med* 2002 ; 136 : 493-503.
- Wielenga RP, Huisveld IA, Bol E, et al. Safety and effects of physical training in chronic heart failure. Results of the chronic heart failure and graded exercise study (CHANGE) *Eur Heart J* 1999 ; 20 : 872-9
- Willenheimer R, Rydberg E, Cline C, et al. Effects on quality of life, symptoms and daily activity 6 months after termination of an exercise training programme in heart failure patients. *Int J Cardiol* 2001 ; 77 : 25-31.

APPAREIL RESPIRATOIRE

- Ambrosino N, Strambi S. New strategies to improve exercise tolerance in chronic obstructive pulmonary disease. *Eur Respir J* 2004 ; 24 : 313-22.
- Beghin L, Michaud L, Turck D, Gottrand F. Technique et intérêt de la mesure de la dépense énergétique et de l'activité physique en recherche clinique chez les patients atteints de mucoviscidose. *Arch Ped* 2005 ; 12 : 1139-44.
- Bradley J, Moran F. Physical training for cystic fibrosis. *The Cochrane Database of Systematic Reviews* 2002, Issue 2. Art. No. : CD002768. DOI : 10.1002/14651858.CD002768.
- Bundgaard A, Ingemann-Hansen T, Halkjaer-Kristensen J, et al. Short-term physical training in bronchial asthma. *Br J Dis Chest* 1983 ; 77 : 147-52.
- Glazebrook C, McPherson AC, Macdonald IA, et al. Asthma as a barrier to children's physical activity : implications for body mass index and mental health. *Pediatrics* 2006 ; 118 : 2443-9.
- Heppner PS, Morgan C, Kaplan RM, Ries AL. Regular walking and long-term maintenance of outcomes after pulmonary rehabilitation. *J Cardiopulm Rehabil* 2006 ; 26 : 44-53.
- Lacasse Y, Brosseau L, Milne S, et al. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2002 ; (3) : CD003793.

Lang DM, Butz AM, Duggan AK, Serwint JR. Physical Activity in Urban School-Aged Children With Asthma. *Pediatrics* 2004 ; 113 ; 341-6.

Matsumoto I, Araki H, Tsuda K, et al. Effects of swimming training on aerobic capacity and exercise induced bronchoconstriction in children with bronchial asthma. *Thorax* 1999 ; 54 : 196-201.

Nagarathna R, Nagendra HR. Yoga for bronchial asthma : a controlled study. *Br Med J* 1985 ; 291 : 1077-9.

Neder JA, Nery LE, Silva AC, et al. Short-term effects of aerobic training in the clinical management of moderate to severe asthma in children. *Thorax* 1999 ; 54 : 202-6.

Nixon PA, Orenstein DM, Kelsey SF. Habitual physical activity in children and adolescents with cystic fibrosis. *Med Sci Sports Exerc* 2001 ; 33 : 30-5.

Puhan MA, Schunemann HJ, Frey M, et al. How should COPD patients exercise during respiratory rehabilitation ? Comparison of exercise modalities and intensities to treat skeletal muscle dysfunction. *Thorax* 2005 ; 60 : 367-75.

Ram FS, Robinson SM, Black PN. Effects of physical training in asthma : a systematic review. *Br J Sports Med* 2000 ; 34 : 162-7.

Richards ML, Davies PS, Bell SC. Energy cost of physical activity in cystic fibrosis. *Eur J Clin Nutr* 2001 ; 55 : 690-7.

Rochester CL. Exercise training in chronic obstructive pulmonary disease. *J Rehabil Res Dev* 2003 ; 40 : S59-80.

Salman GF, Mosier MC, Beasley BW, Calkins DR. Rehabilitation for patients with chronic obstructive pulmonary disease : meta-analysis of randomized controlled trials. *J Gen Intern Med* 2003 ; 18 : 213-21.

Satta A. Exercise training in asthma. *J Sports Med Phys Fitness* 2000 ; 40 : 277-83.

Shames RS, Sharek P, Mayer M, et al. Effectiveness of a multicomponent self-management program in at-risk, school-aged children with asthma. *Ann Allergy Asthma Immunol* 2004 ; 92 : 611-8.

Varray A. L, Mercier JG, Prefaut C.G. Individualized training reduces excessive exercise hyperventilation in asthmatics. *Int Rehabil Res* 1995 ; 18 : 297-312.

Welsh L, Kemp JG, Roberts RG. Effects of physical conditioning on children and adolescents with asthma. *Sports Med* 2005 ; 35 : 127-41.

APPAREIL LOCOMOTEUR

American Geriatrics Society Panel on Exercise and Osteoarthritis. Exercise prescription for older adults with osteoarthritis pain : consensus practice recommendations. A supplement to the AGS Clinical Practice Guidelines on the management of chronic pain in older adults. *J Am Geriatr Soc* 2001 ; 49 : 808-23.

Ashworth NL, Chad KE, Harrison EL, et al. Home versus center based physical activity programs in older adults. *The Cochrane Database of Systematic Reviews* 2005, Issue 1. Art. No. : CD004017.pub2. DOI : 10.1002/14651858.CD004017.pub2

Asikainen TM, Suni JH, Pasanen ME, et al. Effect of brisk walking in 1 or 2 daily bouts and moderate resistance training on lower-extremity muscle strength, balance, and walking performance in women who recently went through menopause : a randomized, controlled trial. *Phys Ther* 2006 ; 86 : 912-23.

Bassey EJ, Ramsdale SJ. Weight-bearing exercise and ground reaction forces : a 12-month randomized controlled trial of effects on BMD in healthy postmenopausal women. *Bone* 1995 ; 16 : 469-76.

Bassey EJ, Rockwell MC, Littlewood JJ, Pye DW. Pre- and post-menopausal women have different bone mineral density response to the same high-impact exercise. *J Bone Miner Res* 1998 ; 13 : 1805-13

Berry MJ, Brubaker PH, O'Toole M, et al. Estimation of VO₂ in older individuals with osteoarthritis of the knee and cardiovascular disease. *Med Sci Sports Exerc* 1996 ; 28 : 808-14.

Braith RW, Mills RM, Welsch MA, et al. Resistance exercise training restores BMD in heart transplant recipients. *J Am Coll Cardiol* 1996 ; 28 : 1471-7.

Bravo G, Gauthier P, Roy PM et al. Impact of a 12-month exercise program on the physical and psychological health of osteopenic women. *J Am Geriatr Soc* 1996 ; 44 : 756-62.

Busch AJ, Barber KAR, Overend TJ, et al. Exercise for treating fibromyalgia syndrome. *The Cochrane Database of Systematic Reviews* 2002, Issue 2. Art. No. : CD003786. DOI : 10.1002/14651858.CD003786.

Chan K, Qin L, Lau M, et al. A randomized prospective study of the effects of Tai Chi Chuan exercises on bone mineral density in postmenopausal women. *Arch Phys Med Rehabil* 2004 ; 85 : 717-22.

Cheng S, Sipila S, Taaffe D, et al. Change in bone mass distribution induced by hormone replacement therapy and high-impact physical activity exercise in post-menopausal women. *Bone* 2002 ; 31 : 126-35.

Chu SP, Kelsey JL, Keegan TH, Sternfeld B et al. Risk factors for proximal humerus fracture. *Am J Epidemiol* 2004 ; 160 : 360-7.

Cobb KL, Bachrach LK, Greendale G, et al. Disordered eating, menstrual irregularity, and bone mineral density in female runners. *Med Sci Sports Exerc* 2003 ; 35 : 711-9

De Feo P, Di Loreto C, Lucidi P, et al. Metabolic response to exercise. *J Endocrinol Invest* 2003 ; 26 : 851-4

Duclos M. Effects of physical training on endocrine functions. *Ann Endocrinol* 2001 ; 62 : 19-32.

Ebrahim S, Thompson Pw, Baskaran V, Evans K. Randomised placebo-controlled trial of brisk walking in the prevention of postmenopausal osteoporosis. *Age Ageing* 1997 ; 26 : 253-60.

Ebrahim S, Thompson PW, Baskaran V, Evans K. Randomized placebo-controlled trial of brisk walking in the prevention of postmenopausal osteoporosis. *Age Ageing* 1997 ; 26 : 253-60.

Engelke K, Kemmler W, Lauber D, et al. Exercise maintains bone density at spine and hip EFOPS : a 3-year longitudinal study in early postmenopausal women. *Osteoporos Int* 2006 ; 17 : 133-42.

Ettinger WH Jr, Burns R, Messier SP, et al. A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis. The Fitness Arthritis and Seniors Trial (FAST). *JAMA* 1997 ; 277 : 25-31.

Feskanich D, Willett W, Colditz G. Walking and leisure-time activity and risk of hip fracture in postmenopausal women. *JAMA* 2002 ; 288 : 2300-6.

Gowans SE, deHueck A. Effectiveness of exercise in management of fibromyalgia. *Curr Opin Rheumatol* 2004 ; 16 : 138-42.

Gregg EW, Cauley JA, Seeley DG, et al. Physical activity and osteoporotic fracture risk in older women. Study of Osteoporotic Fractures Research Group. *Ann Intern Med* 1998 ; 129 : 81-8.

Hayden JA, van Tulder MW, Malmivaara A, Koes BW. Exercise therapy for treatment of non-specific low back pain. *Cochrane Database Syst Rev* 2005 Jul 20 ; (3) : CD000335.

Heaney Rp, Abrams S, Dawson-Hughes B, et al. Peak bone mass. *Osteoporos Int* 2000 ; 11 : 985-1009.

Heikkinen J, Kyllonen E, Kurtilla-Matero E, et al. HRT and exercise : effects on bone density, muscle strength and lipid metabolism. A placebo-controlled 2-year prospective trial on two estrogen-progestin regimens in healthy post-menopausal women. *Maturitas* 1997 ; 26 : 139-49.

Heinonen A, Kannus P, Sievanen H, et al. Randomised controlled trial of effect of high-impact exercise on selected risk factors for osteoporotic fractures. *Lancet* 1996 ; 348 : 1343-7.

Heinonen A, Oja P, Sievanen H, et al. Effect of two training regimens on bone mineral density in healthy perimenopausal women : a randomized controlled trial. *J Bone Miner Res* 1998 ; 13 : 483-90.

Humphries B, Newton RU, Bronks R, et al. Effect of exercise intensity on bone density, strength, and calcium turnover in older women. *Med Sci Sports Exerc* 2000 ; 32 : 1043-50.

Iwamoto J, Takeda T, Otani T, Yabe Y. Effect of increased physical activity on BMD in postmenopausal osteoporotic women. *Keio J Med* 1998 ; 47 : 157-61.

Kemmler W, Von Stengel S, Weineck J, et al. Exercise effects on menopausal risk factors of early postmenopausal women : 3-yr erlangen fitness osteoporosis prevention study results. *Med Sci Sports Exerc* 2005 ; 37 : 194-203.

Kerr D, Morton A, Dick I, Prince R. Exercise effects on bone mass in postmenopausal women are site-specific and load-dependent. *J Bone Miner Res* 1996 ; 11 : 218-25.

Kingsley JD, Panton LB, Toole T, et al. The effects of a 12-week strength-training program on strength and functionality in women with fibromyalgia. *Arch Phys Med Rehabil* 2005 ; 86 : 1713-21.

Kohrt WM, Snead DB, Slatopolsky E, Birge SJ. Additive effects of weight-bearing exercise and estrogen on BMD in older women. *J Bone Miner Res* 1995 ; 10 : 1303-11.

Kujala UM, Kaprio J, Kannus P, et al. Physical activity and osteoporotic hip fracture risk in men. *Arch Intern Med* 2000 ; 160 : 705-8.

Liu-Ambrose TY, Khan KM, Eng JJ, et al. The beneficial effects of group-based exercises on fall risk profile and physical activity persist 1 year postintervention in older women with low bone mass : follow-up after withdrawal of exercise. *J Am Geriatr Soc* 2005 ; 53 : 1767-73.

Lord S, Ward JA, Williams P, Zivanovic E. The effects of a community exercise program on fracture risk factors in older women. *Osteoporos Int* 1996 ; 6 : 361-7.

Maddalozzo GF, Snow CM. High intensity resistance training : effects on bone in older men and women. *Calcif Tissue Int* 2000 ; 66 : 399-404.

Malmivaara A, Hakkinen U, Aro T, et al. The treatment of acute low back pain—bed rest, exercises, or ordinary activity ? *N Engl J Med* 1995 ; 332 : 351-5.

Mannerkorpi K, Iversen MD. Physical exercise in fibromyalgia and related syndromes. *Best Pract Res Clin Rheumatol* 2003 ; 17 : 629-47.

McCartney N, Hicks AL, Martin J, Weber CE. Long-term resistance training in the elderly : effects on dynamic strength, exercise capacity, muscle, and bone. *J Gerontol* 1995 ; 50 : B97-104.

McMurdo MET, Mole PA, Paterson CR. Controlled trial of weight-bearing exercise in older women. *Br Med J* 1997 ; 314 : 569.

Messier SP, Loeser RF, Hoover JL, et al. Osteoarthritis of the knee : effects on gait, strength, and flexibility. *Arch Phys Med Rehabil* 1992 ; 73 : 29-36.

Moffett JK, Togerson D, Bell-Syer S, et al. Randomised controlled trial of exercise for low back pain : clinical outcomes, costs, and preferences. *Br Med J* 1999 ; 319 : 279-83.

Nelson ME, Fiatarone MA, Morganti CM, et al. Effects of high-intensity strength training on multiple risk factors for osteoporotic fracture : a randomized controlled trial. *JAMA* 1994 ; 272 : 1909-14.

Nevitt MC, Cummings SR, Stone KL, et al. Risk factors for a first-incident radiographic vertebral fracture in women > or = 65 years of age : the study of osteoporotic fractures. *J Bone Miner Res* 2005 ; 20 : 131-40.

Nordstrom A, Karlsson C, Nyquist F, et al. Bone loss and fracture risk after reduced physical activity. *J Bone Miner Res* 2005 ; 20 : 202-7

Nordstrom A, Olsson T, Nordstrom P. Bone gained from physical activity and lost through detraining : a longitudinal study in young males. *Osteoporos Int* 2005 ; 16 : 835-41.

Preisinger E, Kerschanch-Schindl K, Wober C, et al. The effect of calisthenic home exercises on postmenopausal fractures—a long-term observational study. *Maturitas* 2001 ; 40 : 61-7

Prince ETR, Devine A, Dick I, et al. The effects of calcium supplementation (milk powder or tablets) and exercise on bone density in postmenopausal women. *J Bone Miner Res* 1995 ; 10 : 1068-75.

Rejeski WJ, Morgan T, et al. A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis. The Fitness Arthritis and Seniors Trial (FAST). *JAMA* 1997 ; 277 : 25-31.

Rhodes EC, Martin AD, Taunton JE, et al. Effects of one year of resistance training on the relation between muscular strength and bone density in elderly women. *Br J Sports Med* 2000 ; 34 : 18-22.

Rockwell JC, Sorensen AM, Baker S. Weight training decreases vertebral bone density in premenopausal women : a prospective study. *J Endocrinol Metabol* 1990 ; 71 : 988-93.

Rogind H, Bibow NB, Jensen B, et al. The effects of a physical training program on patients with osteoarthritis of the knees. *Arch Phys Med Rehabil* 1998 ; 79 : 1421-7.

Rogind H, Bibow-Nielsen B, Jensen B, et al. The effects of a physical training program on patients with osteoarthritis of the knees. *Arch Phys Med Rehabil* 1998 ; 79 : 1421-7.

Salamone LM, Cauley JA, Black DM, et al. Effect of a lifestyle intervention on BMD in premenopausal women : a randomized trial. *Am J Clin Nutr* 1999 ; 70 : 97-103.

Sanchis-Moysi J, Dorado C, Vicente-Rodriguez G, et al. Inter-arm asymmetry in bone mineral content and bone area in postmenopausal recreational tennis players. *Maturitas* 2004 ; 48 : 289-98

Sinaki M, Itoi E, Wahner HW. Stronger back muscles reduce the incidence of vertebral fractures : a prospective 10 year follow-up of postmenopausal women. *Bone* 2002 ; 30 : 836-41.

Tak E, Staats P, Van Hespren A, Hopman-Rock M. The effects of an exercise program for older adults with osteoarthritis of the hip. *J Rheumatol* 2005 ; 32 : 1106-13.

Talbot LA, Gaines JM, Huynh TN, Metter EJ. A home-based pedometer-driven walking program to increase physical activity in older adults with osteoarthritis of the knee : a preliminary study. *J Am Geriatr Soc* 2003 ; 51 : 387-92.

Tsuzuku S, Shimokata H, Ikegami Y, et al. Effects of high versus low-intensity resistance training on bone mineral density in young males. *Calcif Tissue Int* 2001 ; 68 : 342-7

Uusi-Rasi K, Kannus P, Cheng S, et al. Effect of alendronate and exercise on bone and physical performance of postmenopausal women : a randomized controlled trial. *Bone* 2003 ; 33 : 132-43

Vainionpaa A, Korpelainen R, Leppaluoto J, Jamsa T. Effects of high-impact exercise on bone mineral density : a randomized controlled trial in premenopausal women. *Osteoporos Int* 2005 ; 16 : 191-7.

Van Baar M, Assendelft WJ, Dekker J, et al. Effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee : a systematic review of randomized clinical trials. *Arthritis Rheum* 1999 ; 42 : 1361-9

Vico L, Bourrin S, Chatard JC, et al. Possible nonlinear effects of exercise on bone in male subjects over age 60 years. *Anat Rec* 1993 ; 235 : 206-14.

Yamazaki S, Ichimura S, Iwamoto J, et al. Effect of walking exercise on bone metabolism in postmenopausal women with osteopenia/osteoporosis. *J Bone Miner Metab* 2004 ; 22 : 500-8.

NEUROLOGIE

Brown TR, Kraft GH. Exercise and rehabilitation for individuals with multiple sclerosis. *Phys Med Rehabil Clin N Am* 2005 ; 16 : 513-55

Gordon NF, Gulanick M, Costa F, et al. Physical activity and exercise recommendations for stroke survivors : an American Heart Association scientific statement from the Council on Clinical Cardiology, Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention ; the Council on Cardiovascular Nursing ; the Council on Nutrition, Physical Activity, and Metabolism ; and the Stroke Council. *Stroke* 2004 ; 35 : 1230-40.

Heesen C, Romberg A, Gold S, Schulz KH. Physical exercise in multiple sclerosis : supportive care or a putative disease-modifying treatment. *Expert Rev Neurother* 2006 ; 6 : 347-55.

Killeff J, Ashburn A. A pilot study of the effect of aerobic exercise on people with moderate disability multiple sclerosis. *Clin Rehabil* 2005 ; 19 : 165-9.

Leroux A. Exercise training to improve motor performance in chronic stroke : effects of a community-based exercise program. *Int J Rehabil Res* 2005 ; 28 : 17-23.

Macko RF, Ivey FM, Forrester LW, et al. Treadmill exercise rehabilitation improves ambulatory function and cardiovascular fitness in patients with chronic stroke : a randomized, controlled trial. *Stroke* 2005 ; 36 : 2206-11.

Pang MY, Eng JJ, Dawson AS, Gylfadottir S. The use of aerobic exercise training in improving aerobic capacity in individuals with stroke : a meta-analysis. *Clin Rehabil* 2006 ; 20 : 97-111.

Rietberg MB, Brooks D, Uitdehaag BMJ, Kwakkel G. Exercise therapy for multiple sclerosis. *The Cochrane Database of Systematic Reviews* 2004, Issue 3. Art. No. : CD003980. DOI : 10.1002/14651858.CD003980.

Romberg A, Virtanen A, Ruutiainen J. Long-term exercise improves functional impairment but not quality of life in multiple sclerosis. *J Neurol* 2005 ; 252 : 839-45.

Saunders DH, Greig CA, Young A, Mead GE. Physical fitness training for stroke patients. *The Cochrane Database of Systematic Reviews* 2004, Issue 1. Art. No. : CD003316. DOI : 10.1002/14651858.CD003316

CANCÉROLOGIE

Adamsen L, Quist M, Midtgaard J, et al. The effect of a multidimensional exercise intervention on physical capacity, well-being and quality of life in cancer patients undergoing chemotherapy. *Support Care Cancer* 2006 ; 14 : 116-27.

Courneya K, Mackey J, Bell G, et al. Randomized controlled trial of exercise training in postmenopausal breast cancer survivors : Cardiopulmonary and quality of life outcomes. *J Clin Oncol* 2003 ; 21 : 1660-8.

Dimeo F, Fetscher S, Lange W, et al. Effects of aerobic exercise on the physical performance and incidence of treatment-related complications after high-dose chemotherapy. *Blood* 1997 ; 90 : 3390-4.

Dimeo F, Tilmann M, Bertz H, et al. Aerobic exercise in the rehabilitation of cancer patients after high dose chemotherapy and autologous peripheral stem cell transplantation. *Cancer* 1997 ; 79 : 1717-2.

Dimeo FC, Stieglitz RD, Novelli-Fischer U, et al. Effects of physical activity on the fatigue and psychologic status of cancer patients during chemotherapy. *Cancer* 1999 ; 85 : 2273-7.

Evolution de l'incidence et de la mortalité par cancer en France de 1978 à 2000. Réseau français des registres du cancer, Inserm, InVS, 2003.

Frienderreich CM, Courneya KS. Exercise as rehabilitation for cancer patients. *Clin J Sport Med* 1996 ; 6 : 237-44.

Galvao DA, Newton RU. Review of exercise intervention studies in cancer patients. *J Clin Oncol* 2005 ; 23 : 899-909.

Hayes S, Davies PS, Parker T, et al. Quality of life changes following peripheral blood stem cell transplantation and participation in a mixed-type, moderate-intensity, exercise program. *Bone Marrow Transplant* 2004 ; 33 : 553-8.

Hayes S, Davies PS, Parker T, et al. Total energy expenditure and body composition changes following peripheral blood stem cell transplantation and participation in an exercise programme. *Bone Marrow Transplant* 2003 ; 31 : 331-8.

Hayes SC, Davies PS, Parker TW, et al. Role of a mixed type, moderate intensity exercise programme after peripheral blood stem cell transplantation. *Br J Sports Med* 2004 ; 38 : 304-9.

Humpel N, Iverson DC. Review and critique of the quality of exercise recommendations for cancer patients and survivors. *Support Care Cancer* 2005 ; 13 : 493-502.

Hutnick NA, Williams NI, Kraemer WJ, et al. Exercise and lymphocyte activation following chemotherapy for breast cancer. *Med Sci Sports Exerc* 2005 ; 37 : 1827-35.

Jones LW, Eves ND, Courneya KS, et al. Effects of exercise training on antitumor efficacy of doxorubicin in MDA-MB-231 breast cancer xenografts. *Clin Cancer Res* 2005 ; 11 : 6695-8.

Knols R, Aaronson NK, Uebelhart D, et al. Physical exercise in cancer patients during and after medical treatment : a systematic review of randomized and controlled clinical trials. *J Clin Oncol* 2005 ; 23 : 3830-42.

Mock V, Pickett M, Ropka M, et al. Fatigue and quality of life outcomes of exercise during cancer treatment. *Cancer Pract* 2002 ; 9 : 119-27.

Mustian KM, Katula JA, Zhao H. A pilot study to assess the influence of tai chi chuan on functional capacity among breast cancer survivors. *J Support Oncol* 2006 ; 4 : 139-45.

Ness KK, Mertens AC, Hudson MM, et al. Limitations on physical performance and daily activities among long-term survivors of childhood cancer. *Ann Intern Med* 2005 ; 143 : 639-47

Thorsen L, Skovlund E, Stromme SB, et al. Effectiveness of physical activity on cardiorespiratory fitness and health-related quality of life in young and middle-aged cancer patients shortly after chemotherapy. *J Clin Oncol* 2005 ; 23 : 2378-88.

Young-Moo N, Min-Young K, Young-Kyumn K, et al. Exercise therapy effect on natural killer cell cytotoxic activity in stomach cancer patients after curative surgery. *Arch Phys Med Rehabil* 2000 ; 81 : 777-9.

SANTÉ MENTALE

Barbour KA, Blumenthal JA. Exercise training and depression in older adults. *Neurobiol Aging* 2005 ; 26 : 119-23.

Becuwe S, Kalinova E, Le Cren F. Les effets de l'activité physique sur l'estime de soi des personnes de 55 ans et plus. *Vie Vieillesse* 2005 ; 4 : 10-5.

Binsinger C, Laure P, Ambard MF. Regular extra curricular sports practice does not prevent moderate or severe variations in self-esteem or trait anxiety in early adolescents. *J Sports Sci Med* 2006 ; 5 : 123-9.

Blumenthal JA, Babyak MA, Moore KA, et al. Effects of exercise training on older patients with major depression. *Arch Intern Med* 1999 ; 159 : 2349-56.

Bruchon-Schweitzer M, Quintard B. *Personnalité et maladies*. Paris : Dunod, 2001.

Caouette M, Masciotra D, Bouvier H, Bérard J.-M. L'autorégulation de l'anxiété au moyen des activités physiques d'aventure en adopsychiatrie. *Rev Psychoéduc Orientation* 2002 ; 31 : 35-53.

Carmack CL, Boudreaux E, Amaral-Melendez M, et al. Aerobic fitness and leisure physical activity as moderators of the stress-illness relation. *Ann Behav Med* 1999 ; 3 : 251-7.

Coopersmith S. *Self-esteem inventories*. Palo Alto : Consulting Psychologists Press ; 1987.

Dunn AL, Trivedi MH, Kampert JB, et al. Exercise treatment for depression : efficacy and dose response. *Am J Prev Med* 2005 ; 28 : 1-8.

Dunn AL, Trivedi MH, O'Neal HA. Physical activity dose-response effects on outcomes of depression and anxiety. *Med Sci Sports Exerc* 2001 ; 6 : S587-97.

Ekeland E, Heian F, Hagen KB. Can exercise improve self-esteem in children and young people ? A systematic review of randomised controlled trials. *Br J Sports Med* 2005 ; 39 : 792-79.

- Gruber JJ. Physical activity and self-esteem development in children : a meta-analysis. In : Stull GA, Eckert HM (Eds). *Effects of physical activity on children*. Champaign : Human Kinetics, 1986 : 30-48.
- Katula JA, Blissmer BJ, McAuley E. Exercise intensity and self-efficacy effects on anxiety reduction in healthy, older adults. *Behav Med* 1999 ; 3 : 233-47.
- Martinsen EW, Sandvik L, Kolbjornsrud OB. Aerobic exercise in the treatment of nonpsychotic mental disorders : An exploratory study. *Nord Psykia Tidss* 1989 ; 3 : 521-9.
- McAuley E, Mihalko SL, Bane SM. Acute exercise and anxiety reduction : does the environment matter ? *J Sport Exerc Physiol* 1996 ; 18 : 408-419.
- McAuley E. Physical activity and psychosocial outcomes. In : Bouchard C, Shepard RJ, Stephens T (eds). *Physical activity, fitness and health*. Champaign : Human Kinetics 1994 : 551-68.
- McDonald DG, Hodgdon JA. *Psychological effects of aerobic fitness training*. New York : Springer-Verlag, 1991.
- McGee R, Williams S. Does low self-esteem predict health compromising behaviours among adolescents ? *J Adolesc* 2000 ; 5 : 569-82.
- North TC, McCullagh P, VuTran Z. Effect of exercise on depression. *Exerc Sport Sci Rev* 1990 ; 18 : 379-415
- Paluska SA, Schwenk TL. Physical activity and mental health : current concepts. *Sports Med* 2000 ; 29 : 167-80.
- Parfitt G, Eston RG. The relationship between children's habitual activity level and psychological well-being. *Acta Paediatr* 2005 ; 94 : 1791-7.
- Petruzzello S, Landers D, Hatfield B, et al. A meta-analysis on the anxiety-reducing effects of acute and chronic exercise : outcomes and mechanisms. *Sports Med* 1991 ; 11 : 143-82.
- Rosenberg M. *Conceiving the Self*. New-York : Basic, 1979.
- Spence JC, McGannon KR, Poon P. The effect of exercise on global self-esteem : a quantitative review. *Exerc Psychol* 2005 ; 27 : 311-34.
- Spielberger CD. *Manual for the State-Trait Anxiety Inventory*. Palo Alto : Consulting Psychologists Press, 1983.
- Strauss RS, Rodzilsky D, Burack G, Colin M. Psychosocial correlates of physical activity in healthy children. *Arch Pediatr Adolesc Med* 2001 ; 155 : 897-902.
- Wild LG, Flisher AJ, Bhana A, Lombard C. Associations among adolescent risk behaviours and self-esteem in six domains. *J Child Psychol Psych* 2004 ; 8 : 1454-67.

SITUATIONS DE HANDICAP

- Association des Paralysés de France. *Déficiences motrices et situations de handicap : Aspects sociaux, psychologiques, médicaux, techniques, troubles associés*. Paris : APF, 2002.
- Blanc P. *Rapport d'information fait au nom de la commission des affaires sociales sur la politique de compensation du handicap*. N° 369. Sénat, session extraordinaire de 2001-2002.
- Ditor DS, Latimer AE, Ginis KA, et al. Maintenance of exercise participation in individuals with spinal cord injury : effects on quality of life, stress and pain. *Spinal Cord* 2003 ; 41 : 446-50.
- Fragala-Pinkham MA, Haley SM, Rabin J, Kharasch VS. A fitness program for children with disabilities. *Phys Ther* 2005 ; 85 : 1182-200.
- Jacobs PL, Nash MS. Exercise recommendations for individuals with spinal cord injury. *Sports Med* 2004 ; 34 : 727-51.
- Keysor JJ. Does late-life physical activity or exercise prevent or minimize disablement ? A critical review of the scientific evidence. *Am J Prev Med* 2003 ; 3 (S2) : 129-36.
- Kilmer DD, Wright NC, Aitkens S. Impact of a home-based activity and dietary intervention in people with slowly progressive neuromuscular diseases. *Arch Phys Med Rehabil* 2005 ; 86 : 2150-6
- Lancioni GE, Gigante A, O'Reilly MF. Indoor travel and simple tasks as physical exercise for people with profound multiple disabilities. *Percept Mot Skills* 2000 ; 91 : 211-6.
- Mormiche P. Le handicap se conjugue au pluriel. *Insee Première* 2000 ; 742 : 1-4.
- Myslinski MJ. Evidence-based exercise prescription for individuals with spinal cord injury. *J Neurol Phys Ther* 2005 ; 2 : 104-6.
- Nash MS. Exercise as a health-promoting activity following spinal cord injury. *J Neurol Phys Ther* 2005 ; 29 : 87-103, 106.

Raynaud P, Weber A. Les adultes ayant des incapacités lourdes dans les actes essentiels de la vie quotidienne. *DRESS Etudes & Résultats* 2005 ; 439 : 1-11.

Santiago M, Coyle C. Leisure-time physical activity and secondary conditions in women with physical disabilities. *Disabil Rehabil* 2004 ; 26 : 485-94.

Topp R, Boardley D, Morgan AL, et al. Exercise and functional tasks among adults who are functionally limited. *West J Nurs Res* 2005 ; 27 : 252-70.

AUTRES PATHOLOGIES, AUTRES SITUATIONS

Birk TJ, MacArthur RD, Lipton LM, Levine SD. Aerobic exercise training fails to lower hypertriglyceridemia levels in persons with advanced HIV-1 infection. *J Assoc Nurses AIDS Care* 2002 ; 13 : 20-4.

Ciccolo JT, Jowers EM, Bartholomew JB. The benefits of exercise training for quality of life in HIV/AIDS in the post-HAART era. *Sports Med* 2004 ; 34 : 487-99.

Cotman CW, Berchtold NC. Exercise : a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci* 2002 ; 25 : 295-301.

Daniels K, Thornton EW. An analysis of the relationship between hostility and training in the martial arts. *J Sports Sci* 1990 ; 8 : 95-101.

Delva-Tauiiili J. Does brief Aikido training reduce aggression of youth ? *Percep Mot Skills* 1995 ; 80 : 297-8.

Dudgeon WD, Phillips KD, Bopp CM, Hand GA. Physiological and psychological effects of exercise interventions in HIV disease. *AIDS Patient Care STDS* 2004 ; 18 : 81-98.

Dwyer T, Sallis JF, Blizzard L, et al. Relationship of academic performance to physical activity and fitness in children. *Pediatr Exerc Sci* 2001 ; 13 : 225-37.

Edmonds M, McGuire H, Price J. Exercise therapy for chronic fatigue syndrome. *The Cochrane Database of Systematic Reviews* 2004, Issue 3. Art. No. : CD003200. DOI : 10.1002/14651858.CD003200.

Eller LS, Corless I, Bunch EH, et al. Self-care strategies for depressive symptoms in people with HIV disease. *J Adv Nurs* 2005 ; 51 : 119-30.

Field T, Diego M, Sanders CE. Exercise is positively related to adolescents' relationships and academics. *Adolescence* 2001 ; 36 : 105-10.

Fulcher KY, White PD. Randomised controlled trial of graded exercise in patients with chronic fatigue syndrome. *Br Med J* 1997 ; 314 : 1647-52.

Garnier S, Gasnier Y, Mejean L. Impact de l'éducation physique et du petit déjeuner sur les performances académiques des élèves. *Cah Nutr Diet* 2000 ; 35 : 341-7.

Karmisholt K, Gotzsche PC. Physical activity for secondary prevention of disease. Systematic reviews of randomised clinical trials. *Dan Med Bull* 2005 ; 52 : 90-4.

Kim HY, Frongillo EA, Han SS, et al. Academic performance of Korean children is associated with dietary behaviours and physical status. *Asia Pac J Clin Nutr* 2003 ; 12 : 186-92.

Kujala UM. Evidence for exercise therapy in the treatment of chronic disease based on at least three randomized controlled trials—summary of published systematic reviews. *Scand J Med Sci Sports* 2004 ; 14 : 339-45.

Linder KJ. Sport Participation and Perceived Academic Performance of School Children and Youth. *Pediatric Exerc Sci* 1999 ; 11 : 129-44.

Malita FM, Karelis AD, Toma E, Rabasa-Lhoret R. Effects of different types of exercise on body composition and fat distribution in HIV-infected patients : a brief review. *Can J Appl Physiol* 2005 ; 30 : 233-45.

Mior S. Exercise in the treatment of chronic pain. *Clin J Pain* 2001 ; 17 : S77-85.

Moss-Morris R, Sharon C, Tobin R, Baldi JC. A randomized controlled graded exercise trial for chronic fatigue syndrome : outcomes and mechanisms of change. *J Health Psychol* 2005 ; 10 : 245-59.

Mustafa T, Sy FS, Macera CA, et al. Association between exercise and HIV disease progression in a cohort of homosexual men. *Ann Epidemiol* 1999 ; 9 : 127-31.

Nosanchuk TA, MacNeil ML. Examination of the effects of traditional and modern martial arts training on aggressiveness. *Aggressive Behavior* 1989 ; 15 : 153-9.

- Pedersen BK, Saltin B. Evidence for prescribing exercise as therapy in chronic disease. *Scand J Med Sci Sports* 2006 ; 16 : S3-63.
- Reynes E, Lorant J. Competitive martial arts and aggressiveness : a 2-yr. longitudinal study among young boys. *Percept Mot Skills* 2004 ; 98 : 103-15.
- Reynes E, Lorant J. Effect of traditional judo training on aggressiveness among young boys. *Percept Mot Skills* 2002 ; 94 : 21-5.
- Reynolds D, Nicolson RI, Hambly H. Evaluation of an exercise-based treatment for children with reading difficulties. *Dyslexia* 2003 ; 9 : 48-71.
- Rojas R, Hautzinger M, Schlicht W. Effects of exercise training on quality of life, psychological well-being, immune status, and cardiopulmonary fitness in an HIV-1 positive population. *J Sport Exerc Psychol* 2003 ; 25 : 440-55.
- Roubenoff R, Wilson IB. Effect of resistance training on self-reported physical functioning in HIV infection. *Med Sci Sports Exerc* 2001 ; 33 : 1811-7.
- Shephard RJ, Volle M, Lavallee H, et al. Required physical activity and academic grades : a controlled study. In : Ilmarinen J, Valimaki I, eds. *Children and sport*. Berlin : Springer-Verlag ; 1984 : 58-63.
- Shephard RJ. Curricular physical activity and academic performance. *Pediatr Exerc Sci* 1997 ; 9 : 113-25.
- Sibley BA, Etnier JL. The relationship between physical activity and cognition in children : a meta-analysis. *Pediatr Exerc Sci* 2003 ; 15 : 243-56.
- Simmonds MJ, Novy D, Sandoval R. The differential influence of pain and fatigue on physical performance and health status in ambulatory patients with human immunodeficiency virus. *Clin J Pain* 2005 ; 21 : 200-6.
- Smidt N, de Vet HC, Bouter LM, et al. Effectiveness of exercise therapy : a best-evidence summary of systematic reviews. *Aust J Physiother* 2005 ; 51 : 71-85.
- Smith WR, White PD, Buchwald D. A case control study of premorbid and currently reported physical activity levels in chronic fatigue syndrome. *BMC Psychiatry* 2006 ; 6 : 53.
- Taras H. Physical activity and student performance at school. *J Sch Health* 2005 ; 75 : 214-8.
- Tompson PD. Cognitive and behavioral responses to acute exercise in youths : a review. *Pediatr Exerc Sci* 2003 ; 15 : 348-59.
- Trulson ME. Martial arts training : a novel «cure» for juvenile delinquency. *Hum Relations* 1986 ; 39 : 1131-40.
- Yarasheski KE, Roubenoff R. Exercise treatment for HIV-associated metabolic and anthropomorphic complications. *Exerc Sport Sci Rev* 2001 ; 29 : 170-4.
- Zivin G, Hassan NR, DePaula GF, et al. An effective approach to violence prevention : traditional martial arts in middle school. *Adolescence* 2001 ; 36 : 443-59.



4 Prévention et activité physique

PRÉVENTION DES PATHOLOGIES

- Albanes D, Blair A, Taylor PR. Physical activity and risk of cancer in the NHANES I population. *Am J Public Health* 1989 ; 79 : 744-50.
- Andersen LB, Harro M, Sardinha LB et al. Physical activity and clustered cardiovascular risk in children : a cross-sectional study (The European Youth Heart Study). *Lancet* 2006 ; 368 : 299-304.
- Anderson JP, Ross JA, Folsom AR. Anthropometric variables, physical activity, and incidence of ovarian cancer : The Iowa Women's Health Study. *Cancer* 2004 ; 100 : 1515-21.
- Arbman G, Axelson O, Fredriksson M, et al. Do occupational factors influence the risk of colon and rectal cancer in different ways ? *Cancer* 1993 ; 72 : 2543-9.
- Arvers P, Choquet M. Pratiques sportives et consommation de substances psychoactives. Analyse réalisée à partir des données de l'enquête ESPAD 99. *Ann Med Interne* 2003 ; 154 : S25-34.

- Asikainen TM, Kukkonen-Harjula K, Miilunpalo S. Exercise for health for early postmenopausal women : a systematic review of randomised controlled trials. *Sports Med* 2004 ; 34 : 753-78.
- Augestad LB, Schei B, Forsmo S, et al. Healthy postmenopausal women- physical activity and forearm bone mineral density : the Nord-Trondelag health survey. *J Women Aging* 2006 ; 18 : 21-40.
- Bak H, Christensen J, Thomsen BL, et al. Physical activity and risk for lung cancer in a Danish cohort. *Int J Cancer* 2005 ; 116 : 439-44.
- Ballard-Barbash R, Schatzkin A, Albanes D, et al. Physical activity and risk of large bowel cancer in the Framingham Study. *Cancer Res* 1990 ; 50 : 3610-3.
- Bensimhon DR, Kraus WE, Donahue MP. Obesity and physical activity : a review. *Am Heart J* 2006 ; 151 : 598-603.
- Berard A, Bravo G, Gauthier P. Meta-analysis of the effectiveness of physical activity for the prevention of bone loss in postmenopausal women. *Osteoporos Int* 1997 ; 7 : 331-7.
- Bergamaschi A, Morri M, Resi D, et al. Tobacco consumption and sports participation : a survey among university students in northern Italy. *Ann Ig* 2002 ; 14 : 435-42.
- Berlin JA, Colditz GA. A meta-analysis of physical activity in the prevention of coronary heart disease. *Am J Epidemiol* 1990 ; 132 : 612-28.
- Bernstein L, Patel AV, Ursin G, et al. Lifetime recreational exercise activity and breast cancer risk among black women and white women. *J Natl Cancer Inst* 2005 ; 97 : 1671-9.
- Blair SN. Physical activity, fitness, and coronary heart disease. In : Bouchard C, Shephard RJ, Stephens T, editors. *Physical activity, fitness, and health : international proceedings and consensus statement*. Champaign : Human Kinetics, 1994 : 579-90.
- Borer KT. Physical activity in the prevention and amelioration of osteoporosis in women : interaction of mechanical, hormonal and dietary factors. *Sports Med* 2005 ; 35 : 779-830.
- Bradney M, Pearce G, Naughton G, et al. Moderate exercise during growth in prepubertal boys : changes in bone mass, size, volumetric density, and bone strength : a controlled prospective study. *J Bone Miner Res* 1998 ; 13 : 1814-21.
- Broe GA, Creasey H, Jorm AF, et al. Health habits and risk of cognitive impairment and dementia in old age : a prospective study on the effects of exercise, smoking and alcohol consumption. *Aust N Z J Public Health* 1998 ; 22 : 621-3.
- Broe GA, Henderson AS, Creasey H, et al. A case-control study of Alzheimer's disease in Australia. *Neurology* 1990 ; 40 : 1698-707.
- Brown PJ, Josse RG. Clinical practice guidelines for the diagnosis and management of osteoporosis in Canada *CMAJ* 2002 ; 167, 1-34.
- Brownson RC, Chang JC, Davis JR, Smith CA. Physical activity on the job and cancer in Missouri. *Am J Public Health* 1991 ; 81 : 639-42.
- Butler CL, Williams MA, Sorensen TK, et al. Relation between maternal recreational physical activity and plasma lipids in early pregnancy. *Am J Epidemiol* 2004 ; 160 : 350-9.
- Calton BA, Lacey JV Jr, Schatzkin A, et al. Physical activity and the risk of colon cancer among women : a prospective cohort study (United States). *Int J Cancer* 2006 ; 119 : 385-91.
- Cassell C, Benedict M, Specker B. Bone mineral density in elite 7- to 9-yr-old female gymnasts and swimmers. *Med Sci Sports Exerc* 1996 ; 28 : 1243-6.
- Cerhan J, Chiu B, Wallace R, et al. Systolic blood pressure, physical activity, and risk of breast cancer in elderly women. *Am J Epidemiol* 1996 ; 143 : S71.
- Chao A, Connel CJ, Jacobs EJ, et al. Amount, type, and timing of recreational physical activity in relation to colon and rectal cancer in older adults : the Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiol Biomarkers Prev* 2004 ; 13 : 2187-95.
- Chen H, Zhang SM, Schwarzschild MA, et al. Physical activity and the risk of Parkinson disease. *Neurology* 2005 ; 64 : 664-9.
- Chen K, Qiu JL, Zhang Y, Zhao YW. Meta analysis of risk factors cor colorectal cancer. *World J Gastroenterol* 2003 ; 9 : 1598-600.
- Clapp JF, Lopez B, Harcar-Sevcik R. Neonatal behavioral profile of the offspring of women who continued to exercise regularly throughout pregnancy. *Am J Obstet Gynecol* 1999 ; 180 : 91-4.
- Cléroux J, Feldman RF, Petrella RJ. Recommendations on physical exercise training. *Can Med Assoc* 1999 ; 160 : S21-6.
- Colbert LH, Hartman TJ, Maliha N, et al. Physical activity in relation to cancer of the colon and rectum in a cohort of male smokers. *Cancer Epidemiol Biomarkers Prev* 2001 ; 10 : 265-8.
- Colbert LH, Hartman TJ, Tangrea JA, et al. Physical activity and lung cancer risk in male smokers. *Int J Cancer* 2002 ; 98 : 770-3.

Colbert LH, Lacey JV, Schairer C, et al. Physical activity and risk of endometrial cancer in a prospective cohort study. *Cancer Causes Control* 2003 ; 14 : 559-67.

Colbert H A, Lanza E, Ballard-Barbash R, et al. Adenomatous polyp recurrence and physical activity in the Polyp Prevention Trial (United States). *Cancer Causes Control* 2002 ; 13 : 445-53.

Collingwood TR, Sunderlin J, Reynolds R, Kohl HW 3rd. Physical training as a substance abuse prevention intervention for youth. *J Drug Educ* 2000 ; 30 : 435-51.

Cooper C, Cawley M, Bhalla A, et al. Childhood growth, physical activity, and peak bone mass in women. *J Bone Miner Res* 1995 ; 10 : 940-7.

Courteix D, Lespessailles E, Peres SL, et al. Effect of physical training on bone mineral density in prepubertal girls : a comparative study between impact-loading and non-impact-loading sports. *Osteoporos Int* 1998 ; 8 : 152-8.

Cruz-Vidal M, Costas R Jr, García-Palmieri MR, et al. Factors related to diabetes mellitus in Puerto Rican men. *Diabetes* 1979 ; 28 : 300-7.

D'Avanzo B, Nanni O, La Vecchia C, et al. Physical activity and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 1996 ; 5 : 155-60.

Davies GA, Wolfe LA, Mottola MF, MacKinnon C. L'exercice physique pendant la grossesse et le postpartum. *J Obstet Gynaecol Can* 2003 ; 25 : 523-9.

Dempsey JC, Butler CL, Williams MA. No need for a pregnant pause : physical activity may reduce the occurrence of gestational diabetes mellitus and preeclampsia. *Exerc Sport Sci Rev* 2005 ; 33 : 141-9.

Després JP. Visceral obesity, insulin resistance, and dyslipidemia : contribution of endurance exercise training to the treatment of the plurimetabolic syndrome. *Exerc Sport Sci Rev* 1997 ; 25 : 271-300.

Dewey KG, Lovelady CA, Nommsen-Rivers LA, et al. A randomized study of the effects of aerobic exercise by lactating women on breast-milk volume and composition. *N Engl J Med* 1994 ; 330 : 449-53.

Di Pietro L. Physical activity, body weight, and adiposity : an epidemiologic perspective. *Exerc Sport Sci Rev* 1995 ; 23 : 275-303.

Diabetes Prevention Program Research Group. The Diabetes Prevention Program. Design and methods for a clinical trial in the prevention of type 2 diabetes. *Diabetes Care* 1999 ; 22 : 623-34.

Dorn J, Vena J, Brasure J, et al. Lifetime physical activity and breast cancer risk in pre and postmenopausal women. *Med Sci Sports Exerc* 2003 ; 35 : 278-85.

Drinkwater BL, Chesnut CH. Bone density changes during pregnancy and lactation in active women : a longitudinal study. *Bone Mine* 1991 ; 14 : 153-60.

Dumitrescu RG, Cotarla I. Understanding breast cancer risk. Where do we stand in 2005 ? *J Cell Mol Med* 2005 ; 9 : 208-21.

Duppe H, Gardsell P, Johnell O, Ornstein E. Bone mineral density in female junior, senior and former football players. *Osteoporos Int* 1996 ; 6 : 437-41.

Eriksson J, Lindstrom J, Valle T, et al. Prevention of Type II diabetes in subjects with impaired glucose tolerance : the Diabetes Prevention Study (DPS) in Finland. Study design and 1-year interim report on the feasibility of the lifestyle intervention programme *Diabetologia* 1999 ; 42 : 793-801.

Eriksson KF, Lindgarde F. Prevention of type 2 (non-insulin-dependent) diabetes mellitus by diet and physical exercise. The 6-year Malmo feasibility study. *Diabetologia* 1991 ; 34 : 891-8.

Favier F, Fianu A, Naty N, et al. Essai de prévention primaire du diabète de type 2 et du syndrome métabolique à la Réunion. *Rev Med Assur Maladie* 2005 ; 36 : 5-13.

Florack EI, Zielhuis GA, Rolland R. The influence of occupational physical activity on the menstrual cycle and fecundability. *Epidemiology* 1994 ; 5 : 14-8.

Folsom AR, Prineas RJ, Kaye SA, Munger RG. Incidence of hypertension and stroke in relation to body fat distribution and other risks factors in older women. *Stroke* 1990 ; 21 : 701-6.

Fransson E, de Faire U, Ahlbom A, et al. The effect of leisure-time physical activity on the risk of acute myocardial infarction depending on body mass index : a population-based case-control study. *BMC Public Health* 2006 ; 6 : 296.

Fredriksson M, Bengtsson NO, Hardell L, Axelson O. Colon cancer, physical activity, and occupational exposures. A case-control study. *Cancer* 1989 ; 63 : 1838-42.

French SA, Jeffery RW, Forster JL, et al. Predictors of weight change over two years among a population of working adults : the Healthy Worker Project. *Int J Obes* 1994 ; 18 : 145-54.

Friedenreich CM, McGregor SE, Courneya KS, et al. Casecontrol study of lifetime total physical activity and prostate cancer risk. *Am J Epidemiol* 2004 ; 159 : 740-9.

Friedenreich CM. Physical activity and cancer prevention : from observational to intervention research. *Cancer Epidemiol Biomarkers Prev* 2001 ; 10, 287-301.

Gallagher RP, Huchcroft S, Phillips N, et al. Physical activity, medical history, and risk of testicular cancer. *Cancer Causes Control* 1995 ; 6 : 398-406.

Gammon MD, John EM, Britton JA. Recreational and occupational physical activities and risk of breast cancer. *J Natl Cancer Inst* 1998 ; 90 : 100-17.

Gerhardsson M, Norell SE, Kiviranta H, et al. Sedentary jobs and colon cancer. *Am J Epidemiol* 1986 ; 123 : 775-80.

Gerhardsson M, Steineck G, Hagman U, et al. Physical activity and colon cancer : a case-referent study in Stockholm. *Int J Cancer* 1990 ; 46 : 985-9.

Gillespie LD, Gillespie WJ, Robertson MC, et al. Interventions for preventing falls in elderly people. *Cochrane Database Syst Rev* 2003 ; (4) : CD000340.

Giovannucci E, Ascherio A, Rimm EB, et al. Physical activity, obesity, and risk for colon cancer and adenoma in men. *Ann Int Med* 1995 ; 122 : 327-34.

Glazener CMA, Herbison GP, Wilson PD, et al. Conservative management of persistent postnatal urinary and faecal incontinence : Randomised controlled trial. *Br Med J* 2001 ; 323 : 593-6.

Hambrecht R, Niebauer J, Marburger C, et al. V Various intensities of leisure time activity in patients with coronary artery arious disease : effects on cardiorespiratory fitness and progression of coronary ather- atherosclerotic lesions. osclerotic J Am Coll Cardiol 1993 ; 22 : 468-77.

Hanley AJ, Johnson KC, Villeneuve PJ, Mao Y. Physical activity, anthropometric factors and risk of pancreatic cancer : results from the Canadian enhanced cancer surveillance system. *Int J Cancer* 2001 ; 94 : 140-7.

Hannan LM, Leitzmann MF, Lacey JV Jr, et al. Physical activity and risk of ovarian cancer : a prospective study in the United States. *Cancer Epidemiol Biomarkers Prev* 2004 ; 13 : 765-70

Harber VJ. Energy balance and reproductive function in active women. *Can J Appl Physiol* 2003 ; 29 : 48-58.

Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS Jr. Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. *N Engl J Med* 1991 ; 325 : 147-52.

Hetland ML, Haarbo J, Christiansen C. Low bone mass and high bone turnover in male long distance runners. *J Clin Endocrinol Metab* 1993 ; 77 : 770-5.

Higgins S, Gill JMR, Janilyonite R, et al. Physical activity, dietary intake and metabolic risk factors in non-diabetic daughters of patients with type II diabetes. *Prev Med* 2005 ; 40 : 140-51.

Hill JO, Wyatt HR, Reed GW, Peters JC. Obesity and the environment : Where do we go from here ? *Science* 2003 ; 299 : 853-5

Hjollund NH, Storgaard L, Ernst E, et al. The relation between daily activities and scrotal temperature. *Reprod Toxicol* 2002 ; 16 : 209-14.

Hsiung GY, Sadovnick AD, Feldman H. Apolipoprotein E epsilon4 genotype as a risk factor for cognitive decline and dementia : data from the Canadian Study of Health and Aging. *CMAJ* 2004 ; 171 : 863-7.

Hu FB, Manson JE, Stampfer MJ, et al. Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *N Engl J Med* 2001 ; 345 : 790-7.

Hu FB, Sigal RJ, Rich-Edwards JW, et al. Walking compared with vigorous physical activity and risk of type 2 diabetes in women : a prospective study. *JAMA* 1999 ; 282 : 1433-9.

Hu G, Qiao Q, Silventoinen K, et al. Occupational, commuting, and leisure-time physical activity in relation to risk for type 2 diabetes in middle-aged Finnish men and women. *Diabetologia* 2003 ; 46 : 322-9.

Janz KF, Gilmore JM, Burns TL, et al. Physical activity augments bone mineral accrual in young children : The Iowa Bone Development study. *J Pediatr* 2006 ; 148 : 793-9.

Jian L, Shen ZJ, Lee HA, Binns CW. Moderate physical activity and prostate cancer risk : a case-control study in China. *Eur J Epidemiol* 2005 ; 20 : 155-60.

Jung A, Eberl M, Schill WB. Improvement of semen quality by nocturnal scrotal cooling and moderate behavioural change to reduce genital heat stress in men with oligoasthenoteratozoospermia. *Reproduction*. 2001 ; 121 : 595-603

Kardel KR, Kase T. Training in pregnant women : effects on fetal development and birth. *Am J Obstet Gynecol* 1998 ; 178 : 280-6.

Karlsson MK, Ahlborg HG, Karlsson C. Maternity and bone mineral density. *Acta Orthop* 2005 ; 76 : 2-13.

Kato I, Tominaga S, Ikari A. A case-control study of male colorectal cancer in Aichi Prefecture, Japan : with special reference to occupational activity level, drinking habits and family history. *Jpn J Cancer Res* 1990 ; 81 : 115-21

Kawate R, Yamakido M, Nishimoto Y, et al. Diabetes mellitus and its vascular complications in Japanese migrants on the island of Hawaii. *Diabetes Care* 1979 ; 2 : 161-70.

Keller G, Tran ZV. Aerobic exercise and normotensive adults : a meta-analysis. *Med Sci Sport Exerc.* 1995 ; 27 : 1371-7

Kemmeler W, Weineck J, Kalender WA, Engelke K. The effect of habitual physical activity, non-athletic exercise, muscle strength, and VO2max on bone mineral density is rather low in early postmenopausal osteopenic women. *J Musculoskelet Neuronal Interact* 2004 ; 4 : 325-34.

King H, Kriska AM. Prevention of type II diabetes by physical training : epidemiology considerations and study methods. *Diabetes Care* 1992 ; 15 : 1794-9.

Kochan-Vintinner A. *Active Living During Pregnancy : Physical Activity Guidelines for Mother and Baby*. Ottawa : Canadian Society for Exercise Physiology and Health ; 1999.

Kohrt WM, Bloomfield SA, Little KD, et al. American College of Sports Medicine Position Stand : physical activity and bone health. *Med Sci Sports Exerc.* 2004 ; 11 : 1985-1996.

Kontulainen S, Kannus P, Haapasalo H, et al. Changes in bone mineral content with decreased training in competitive young adult tennis players and controls : A prospective 4-yr follow-up. *Med Sci Sports Exerc* 1999 ; 31 : 646-52.

Kosaka K, Noda M, Kuzuya T. Prevention of type 2 diabetes by lifestyle intervention : a Japanese trial in IGT males. *Diabetes Res Clin Pract* 2005 ; 67 : 152-62.

Kramer MS. Aerobic exercise for women during pregnancy. *The Cochrane Database of Systematic Reviews* 2002, Issue 2. Art. No. : CD000180. DOI : 10.1002/14651858.CD000180.

Kriska AM, Blair SN, Pereira MA. The potential role of physical activity in the prevention of noninsulindependent diabetes mellitus : the epidemiological evidence. *Exerc Sport Sci Rev* 1994 ; 22 : 121-43.

Kristal AR, Littman AJ, Benitez D, White E. Yoga practice is associated with attenuated weight gain in healthy, middle-aged men and women. *Altern Ther Health Med* 2005 ; 11 : 28-33.

Kruk J, Aboul-Enein HY. Physical activity in the prevention of cancer. *Asian Pac J Cancer Prev* 2006 ; 7 : 11-21.

Kune GA, Kune S, Watson LF. Body weight and physical activity as predictors of colorectal cancer risk. *Nutr Cancer* 1990 ; 13 : 9-17.

Kushi LH, Byers T, Doyle C, et al. American Cancer Society Guidelines on Nutrition and Physical Activity for cancer prevention : reducing the risk of cancer with healthy food choices and physical activity. *CA Cancer J Clin* 2006 ; 56 : 254-81.

Laaksonen DE, Lindstrom J, Lakka TA, et al. Physical activity in the prevention of type 2 diabetes : the Finnish diabetes prevention study. *Diabetes* 2005 ; 54 : 158-65.

Larson EB, Wang L, Bowen JD, et al. Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. *Ann Intern Med* 2006 ; 144 : 73-81.

Larson-Meyer DE. Effect of postpartum exercise on mothers and their offspring : a review of the literature. *Obes Rev* 2002 ; 10 : 841-53.

Laure P, Binsinger C, Picard J. Effets de la gymnastique d'entretien sur la consommation de médicaments psychotropes par les seniors. *Sci Sports* 2005 ; 20 : 297-99.

Laurin DL, Verreault R, Lindsay J, et al. Physical activity and risk of cognitive impairment and dementia in elderly persons. *Arch Neurol* 2001 ; 58 : 498-504.

Lautenschlager NT, Almeida OP. Physical activity and cognition in old age. *Curr Opin Psychiatry* 2006 ; 19 : 190-3.

Lee IM, Manson JE, Ajani U, et al. Physical activity and risk of colon cancer : The Physicians' Health Survey (United States). *Cancer Causes Control* 1997 ; 8 : 568-74.

Lee IM, Paffenbarger RS, Hsieh CC. Physical activity and risk of developing colorectal cancer among college alumni. *J Natl Cancer Inst* 1991 ; 83 : 1324-9.

Leiferman JA, Evenson KR. The effect of regular leisure physical activity on birth outcomes. *Matern Child Health J* 2003 ; 7 : 59-64

Leitzmann MF, Rimm EB, Willett WC, et al. Recreational physical activity and the risk of cholecystectomy in women. *New Engl J Med* 1999 ; 341 : 777-84

- Lerchl A, Keck C, Spiteri-Grech J, Nieschlag E. Diurnal variations in scrotal temperature of normal men and patients with varicocele before and after treatment. *Int J Androl* 1993 ; 16 : 195–200.
- Li G, Shen YC, Chen CH, et al. An epidemiological survey of age-related dementia in an urban area of Beijing. *Acta Psychiatr Scand* 1989 ; 79 : 557-63.
- Lindstrom J, Ilanne-Parikka P, Peltonen M, et al. Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention : follow-up of the Finnish Diabetes Prevention Study. *Lancet* 2006 ; 368 : 1673-9.
- Longnecker MP, Gerhardsson le Verdier M, Frumkin H, Carpenter C. A case-control study of physical activity in relation to risk of cancer of the right colon and rectum in men. *Int J Epidemiol* 1995 ; 24 : 42-50.
- Loucks AB. Energy availability, not body fatness, regulates reproductive function in women. *Exerc Sport Sci Rev* 2003 ; 31 : 144-8.
- Lovelady CA, Garner KE, Moreno KL, Williams JP. The effect of weight loss in overweight, lactating women on the growth of their infant. *N Engl J Med* 2000 ; 342 : 449-53.
- MacAuley D, McCrum EE, Stott G, et al. Physical activity, physical fitness, blood pressure, and fibrinogen in the Northern Ireland health and activity survey. *J Epidemiol Comm Health* 1996 ; 50 : 258-63.
- Macera CA. Past recreational physical activity and risk of breast cancer. *Clin J Sport Med* 2005 ; 15 : 115-6.
- MacKelvie KJ, Petit MA, Khan KM, et al. Bone mass and structure are enhanced following a 2-year randomized controlled trial of exercise in prepubertal boys. *Bone* 2004 ; 34 : 755-64.
- Malmberg J, Miilunpalo S, Pasanen M et al. Characteristics of leisure time physical activity associated with risk of decline in perceived health—a 10-year follow-up of middle-aged and elderly men and women. *Prev Med* 2005 ; 41 : 141-50.
- Manson JE, Rimm EB, Stampfer MJ, et al. Physical activity and incidence of non-insulin-dependent diabetes in women. *Lancet* 1991 ; 338 : 774-8.
- Mao Y, Pan S, Wen SW, et al. Physical inactivity, energy intake, obesity and the risk of rectal cancer in Canada. *Int J Cancer* 2003 ; 105 : 831-7.
- Mao Y, Pan S, Wen SW, Johnson KC. The Canadian Cancer. Physical activity and the risk of lung cancer in Canada. *Am J Epidemiol* 2003 ; 158 : 564-75.
- Marcus PM, Newcomb PA, Storer BE. Early adulthood physical activity and colon cancer risk among Wisconsin women. *Cancer Epidemiol Biomarkers Prev* 1994 ; 3 : 641-4.
- Margolis KL, Mucci L, Braaten T, et al. Physical activity in different periods of life and the risk of breast cancer : the Norwegian-Swedish Women's Lifestyle and Health cohort study. *Cancer Epidemiol Biomarkers Prev* 2005 ; 14 : 27-32.
- Masson JE, Hu FB, Rich-Edwards JW, et al. A prospective study of walking as compared with a rigorous exercise in the prevention of coronary heart disease in women. *N Engl J Med* 1999 ; 341 : 650-58
- Matthews CE, Xu WH, Zheng W, et al. Physical activity and risk of endometrial cancer : a report from the Shanghai endometrial cancer study. *Cancer Epidemiol Biomarkers Prev* 2005 ; 14 : 779-85.
- Mayoux-Benhamou MA, Leyge JF, Roux C, Revel M. Cross-sectional study of weight-bearing activity on proximal femur bone mineral density. *Calcif Tissue Int* 1999 ; 64 : 179-83.
- Meher S, Duley L. Exercise or other physical activity for preventing pre-eclampsia and its complications. *Cochrane Database Syst Rev* 2006 Apr 19 ; (2) : CD005942.
- Menezes RJ, Tomlinson G, Kreiger N. Physical activity and risk of renal cell carcinoma. *Int J Cancer* 2003 ; 107 : 642-6.
- Mertens AJ, Sweeney C, Shahar E, Rosamond WD, Folsom AA. Physical activity and breast cancer incidence in middle-aged women : a prospective cohort study. *Breast Cancer Res Treat* 2005 ; Dec 2 : 1-6.
- Mink PJ, Folsom AR, Sellers TA, Kushi LH. Physical activity, waist-to-hip ratio, and other risk factors for ovarian cancer : a follow-up study of older women. *Epidemiology* 1996 ; 7 : 38-45.
- Mittleman MA, Maclure M, Tofler G, et al. Triggering of acute myocardial infarction by heavy physical exertion. *N Engl J Med* 1993 ; 329 : 1677-83.
- Mogren IM. Previous physical activity decreases the risk of low back pain and pelvic pain during pregnancy. *Scand J Public Health* 2005 ; 33 : 300-6.
- Morkved S, Bo K. Effect of postpartum pelvic floor muscle training in prevention and treatment of urinary incontinence : a one-year follow up. *Br J Obstet Gynaecol* 2000 ; 107 : 1022-8.
- Morris FL, Naughton GA, Gibbs JL, et al. Prospective ten-month exercise intervention in premenarcheal girls : Positive effects on bone and lean mass. *J Bone Miner Res* 1997 ; 12 : 1453-62.

Morris JN, Heady JA, Raffle PAB, Roberts CGQ, Parks JW Coronary heart disease and physical activity of work. *Lancet* 1953 ; 2 : 1111-20.

Nerin I, Crucelaegui A, Novella P, et al. A survey on the relationship between tobacco use and physical exercise among university students. *Arch Bronconeumol* 2004 ; 40 : 5-9.

Nilsen TI, Romundstad PR, Vatten LJ. Recreational physical activity and risk of prostate cancer : A prospective population-based study in Norway (the HUNT study). *Int J Cancer* 2006 ; 119 : 2943-7.

Nordstrom P, Pettersson U, Lorentzon R. Type of physical activity, muscle strength, and pubertal stage as determinants of bone mineral density and bone area in adolescent boys. *J Bone Miner Res* 1998 ; 13 : 1141-8.

Ohtake PJ, Wolfe LA. Physical conditioning attenuates respiratory responses to steady-state exercise in late gestation. *Med Sci Sports Exerc* 1998 ; 30 : 17-27.

Okruhlica L, Kaco J, Klempova D. Sports activities in the prevention of heroin dependency. *Eur Addict Res* 2001 ; 7 : 83-6.

Paffenbarger RS Jr, Hyde RT, et al. The association of changes in physical-activity level and other lifestyle characteristics with mortality among men. *N Engl J Med* 1993 ; 328 : 538-45.

Pan SY, Ugnat AM, Mao Y. Physical activity and the risk of ovarian cancer : a case-control study in Canada. *Int J Cancer* 2005 ; 117 : 300-7.

Pan XR, Li GW, Hu YH, et al. Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. *Diabetes Care* 1997 ; 20 : 537-44.

Patel AV, Rodriguez C, Jacobs EJ, et al. Recreational physical activity and risk of prostate cancer in a large cohort of US men. *Cancer Epidemiol Biomarkers Prev* 2005 ; 14 : 275-9.

Patel AV, Rodriguez C, Pavluck AL, et al. Recreational physical activity and sedentary behavior in relation to ovarian cancer risk in a large cohort of US women. *Am J Epidemiol* 2006 ; 163 : 709-16.

Perrig-Chiello P, Perrig WJ, Ehrsam R, et al. The effects of resistance training on well-being and memory in elderly volunteers. *Age Ageing* 1998 ; 27 : 469-75

Petit MA, McKay HA, MacKelvie KJ, et al. A randomized school-based jumping intervention confers site and maturity-specific benefits on bone structural properties in girls : a hip structural analysis study. *J Bone Miner Res* 2002 ; 17 : 363-72.

Physical Activity and Health : A Report of the Surgeon General. Atlanta, GA : US Department of Health and Human Services, Centers for Disease Control and Prevention, 1996

Poudevigne MS, O'Connor PJ. A review of physical activity patterns in pregnant women and their relationship to psychological health. *Sports Med* 2006 ; 36 : 19-38.

Poudevigne MS, O'Connor PJ. Physical activity and mood during pregnancy. *Med Sci Sports Exerc* 2005 ; 37 : 1374-80.

Pukkala E, Kaprio J, Koskenvuo M, et al. Cancer incidence among Finnish world class male athletes. *Int J Sports Med* 2000 ; 21 : 216-20.

Pukkala E, Poskiparta M, Apter D, Vihko V. Life-long physical activity and cancer risk among Finnish female teachers. *Eur J Cancer Prev* 1993 ; 2 : 369-76.

Qiu D, Kurosawa M, Lin Y, et al. Overview of the epidemiology of pancreatic cancer focusing on the JACC Study. *J Epidemiol* 2005 ; 15S2 : S157-67.

Quinn TJ, Carey GB. Does exercise intensity or diet influence lactic acid accumulation in breast milk ? *Med Sci Sports Exerc* 1999 ; 31 : 105-10.

Rabkin CS, Anderson HR, Bland JM, et al. Maternal activity and birth weight : a prospective, population-based study. *Am J Epidemiol* 1990 ; 131 : 522-31.

Ravussin E, Bennett PH, Valencia ME, et al. Effects of a traditional lifestyle on obesity in Pima Indians. *Diabetes Care* 1994 ; 17 : 1067-74.

Redman LM. Physical activity and its effects on reproduction. *Reprod Biomed Online* 2006 ; 12 : 579-86.

Regan C, Katona C, Walker Z, Livingston G. Relationship of exercise and other risk factors to depression of Alzheimer's disease : the LASER-AD study. *Int J Geriatr Psychiatry* 2005 ; 20 : 261-8.

Renaud M, Bherer L. L'impact de la condition physique sur le vieillissement cognitif. *Psychol Neuropsychiat Vieillesse* 2005 ; 3 : 199-206.

Rich-Edwards JW, Spiegelman D, Garland M, et al. Physical activity, body mass index, and ovulatory disorder infertility. *Epidemiology* 2002 ; 13 : 184-90.

Rockhill B, Willett WC, Hunter DJ, et al. Physical activity and breast cancer risk in a cohort of young women. *N Eng J Med* 1998 ; 90 : 1155-60.

Rossing MA, Remler R, Voigt LF, et al. Recreational physical activity and risk of papillary thyroid cancer. *Cancer Causes Control* 2001 ; 12 : 881-5.

Rovio S, Kareholt I, Helkala EL, et al. Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's disease. *Lancet Neurol* 2005 ; 4 : 705-11.

Rudra CB, Williams MA, Lee IM, et al. Perceived exertion in physical activity and risk of gestational diabetes mellitus. *Epidemiology* 2006 ; 17 : 31-7.

Rundle A. Molecular epidemiology of physical activity and cancer. *Cancer Epidemiol Biomarkers Prev* 2005 ; 14 : 227-36.

Samad AK, Taylor RS, Marshall T, Chapman MA. A meta-analysis of the association of physical activity with reduced risk of colorectal cancer. *Colorectal Dis* 2005 ; 7 : 204-13.

Schneider JE. Energy balance and reproduction. *Physiol Behav* 2004 ; 81 : 289-317.

Schnohr P, Gronbaek M, Petersen L, et al. Physical activity in leisure-time and risk of cancer : 14-year follow-up of 28000 Danish men and women. *Scand J Public Health* 2005 ; 33 : 244-9.

Schroder H, Marrugat J, Elosua R, Maribel I. Relationship between body mass index, serum cholesterol, leisure-time physical activity, and diet in a Mediterranean Southern-Europe population *Br J Nutr* 2003 ; 90, 431-43.

Schroder H, Marrugat J, Elosua R, et al. Relationship between body mass index, serum cholesterol, leisure-time physical activity, and diet in a Mediterranean Southern-Europe population. *Br J Nutr* 2003 ; 90 : 431-9.

Schulz LO, Schoeller DA. A compilation of total daily energy expenditures and body weights in healthy adults. *Am J Clin Nutr* 1994 ; 60 : 676-81.

Severson RK, Nomura AM, Grove JS, Stemmermann GN. A prospective analysis of physical activity and cancer. *Am J Epidemiol* 1989 ; 130 : 522-9

Shors AR, Solomon C, McTiernan A, White E. Melanoma risk in relation to height, weight, and exercise. *Cancer Causes Control* 2001 ; 12 : 599-606.

Simon C. Prévention de l'obésité. *Rev Prat* 2005 ; 55 : 1453-60.

Slattery ML, Edwards S, Curtin K, et al. Physical activity and colorectal cancer. *Am J Epidemiol* 2003 ; 158 : 214-24.

Slattery ML, Jacobs DR Jr, Nichaman MZ. Leisure-time physical activity and coronary heart disease death : the U.S. Railroad Study. *Circulation* 1989 ; 79 : 304-11.

Slattery ML. Physical activity and colorectal cancer. *Sports Med* 2004 ; 34 : 239-52.

Srivastava A, Kreiger N. Relation of physical activity to risk of testicular cancer. *Am J Epidemiol* 2000 ; 151 : 78-87.

Steindorf K, Jedrychowski W, Schmidt M, et al. Case-control study of lifetime occupational and recreational physical activity and risks of colon and rectal cancer. *Eur J Cancer Prev* 2005 ; 14 : 363-71.

Sternfeld B, Quesenberry CP Jr, Eskenazi B, Newman LA. Exercise during pregnancy and pregnancy outcome. *Med Sci Sports Exerc* 1995 ; 27 : 634-40.

Sternfeld B, Quesenberry JR, Eskenazi B, Newman LA. Exercise during pregnancy and pregnancy outcome. *Med Sci Sports Exerc* 1995 ; 27 : 634-40.

Storti KL, Brach JS, FitzGerald SJ, et al. Physical activity and decreased risk of clinical gallstone disease among post-menopausal women. *Prev Med* 2005 ; 41 : 772-7

Stoy J, Hjollund NH, Mortensen JT, et al. Semen quality and sedentary work position. *Int J Androl* 2004 ; 27 : 5-11.

Summerbell CD, Waters E, Edmunds LD et al. Interventions for preventing obesity in children. *Cochrane Database Syst Rev* 2005 Jul 20 ; (3) : CD001871.

Takito MY, Benicio MH, Latorre Mdo R. Maternal posture and its influence on birthweight. *Rev Saude Publica* 2005 ; 39 : 325-32.

Tardon A, Lee WJ, Delgado-Rodriguez M, et al. Leisure-time physical activity and lung cancer : a meta-analysis. *Cancer Causes Control* 2005 ; 16 : 389-97.

Tavani A, Braga C, La Vecchia C, et al. Physical activity and risk of cancers of the colon and rectum : an Italian cas-control study. *Br J Cancer* 1999 ; 79 : 1912-6.

Taylor RJ, Bennett PH, LeGonidec G, et al. The prevalence of diabetes mellitus in a traditional-living Polynesian population : the Wallis Island survey. *Am J Pub Health* 1983 ; 6 : 334-40.

Tehard B, Friedenreich CM, Oppert JM, Clavel-Chapelon F. Effect of Physical Activity on Women at Increased Risk of Breast Cancer : Results from the E3N Cohort Study. *Cancer Epidemiol Biomarkers Prev* 2006 ; 15 : 57-64.

The Diabetes Prevention Program Research Group. The Diabetes Prevention Program. *Diabetes Care* 2000 ; 23 : 1619-29.

Thompson RF, Crist DM, Marsh M, Rosenthal M. Effects of physical exercise for elderly patients with physical impairments. *J Am Geriatr Soc* 1988 ; 36 : 130-5.

Thune I, Lund E. Physical activity and risk of colorectal cancer in men and women. *Br J Cancer* 1996 ; 73 : 1134-40.

Thune I, Lund E. Physical activity and the risk of prostate and testicular cancer : a cohort study of 53,000 Norwegian men. *Cancer Causes Control* 1994 ; 5 : 549-56.

Thune I, Lund E. The influence of physical activity on lung-cancer risk : A prospective study of 81,516 men and women. *Int J Cancer* 1997 ; 70 : 57-62.

Tuomilehto J, Lindstrom J, Eriksson JG, et al. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 2001 ; 344 : 1343-1350.

United Kingdom Testicular Cancer Study Group. Aetiology of testicular cancer : association with congenital abnormalities, age at puberty, infertility, and exercise. *Br Med J* 1994 ; 308 : 1393-9

Ussher M. Exercise interventions for smoking cessation. *Cochrane Database Syst Rev* 2005 Jan 25 ; (1) : CD002295.

Valdimarsson O, Linden C, Johnell O, et al. Daily physical education in the school curriculum in prepubertal girls during 1 year is followed by an increase in bone mineral accrual and bone width—data from the prospective controlled Malmo pediatric osteoporosis prevention study. *Calcif Tissue Int* 2006 ; 78 : 65-71.

van Dijk BA, Schouten LJ, Kiemeneij LA, et al. Relation of height, body mass, energy intake, and physical activity to risk of renal cell carcinoma : results from the Netherlands Cohort Study. *Am J Epidemiol* 2004 ; 160 : 1159-67.

Vena JE, Graham S, Zielezny M, et al. Lifetime occupational exercise and colon cancer. *Am J Epidemiol* 1985 ; 122 : 357-65.

Villeneuve PJ, Johnson KC, Kreiger N, Mao Y. Risk factors for prostate cancer : results from the Canadian National Enhanced Cancer Surveillance System. The Canadian Cancer Registries Epidemiology Research Group. *Cancer Causes Control* 1999 ; 10 : 355-67.

Wagner A, Simon C, Ducimetière P, et al. Influence de la quantité et du type d'activité physique sur l'adiposité et la prise de poids chez des hommes d'âge moyen. *Nutr Clin Metab* 1999 ; 13 : 64S.

Wagner A, Simon C, Evans A, et al. Physical activity event incident in north ireland and france : the prospective study of myocardial infraction (PRIME) *Circulation* 2002 ; 105 : 2247-52.

Wallace BA, Cumming RG. Systematic review of randomized trials of the effect of exercise on bone mass in pre- and postmenopausal women. *Calcif Tissue Int* 2000 ; 67 : 10-8

Wannamethee SG, Shaper AG, Walker M. Physical activity and risk of cancer in middle-aged men. *Br J Cancer* 2001 ; 85 : 1311-6.

Webb KA, Wolfe LA, McGrath MJ. Effects of acute and chronic maternal exercise on fetal heart rate. *J Appl Physiol* 1994 ; 77 : 2207-13.

Wei EK, Giovannucci E, Wu K, et al. Comparison of risk factors for colon and rectal cancer. *Int J Cancer* 2004 ; 108 : 433-42.

Werch C, Moore M, DiClemente CC, et al. A sport-based intervention for preventing alcohol use and promoting physical activity among adolescents. *J Sch Health* 2003 ; 73 : 380-8.

White E, Jacobs EJ, Daling JR. Physical activity in relation to colon cancer in middle-aged men and women. *Am J Epidemiol* 1996 ; 144 : 42-50.

Whittemore A, Wu-Williams AJ, Lee M, et al. Diet, physical activity and colorectal cancer among Chinese in North America and China. *J Natl Cancer Inst* 1990 ; 82 : 915-26.

Wing RR, Venditti E, Jakicic JM, et al. Lifestyle intervention in overweight individuals with a family history of diabetes. *Diabetes Care* 1998 ; 21 : 350-9.

Wolff I, Van Croonenborg JJ, Kemper HC, et al. The effect of exercise training program on bone mass : a meta-analysis of published controlled trials in pre- and post-menopausal women. *Osteoporosis Int* 1999 ; 9 : 1-12

Zang M, Lee AH, Binns CW. Physical activity and epithelial ovarian cancer risk : a case-control study in china. *Int J Cancer* 2003 ; 105 : 838-43.

Zeegers MP, Dirx MJ, van den Brandt PA. Physical activity and the risk of prostate cancer in the Netherlands cohort study, results after 9,3 years of follow-up. *Cancer Epidemiol Biomarkers Prev* 2005 ; 14 : 1490-5.

Zhang C, Solomon CG, Manson JE, Hu FB. A prospective study of pregravid physical activity and sedentary behaviors in relation to the risk for gestational diabetes mellitus. *Arch Intern Med* 2006 ; 166 : 543-8.

Zhang Y, Cantor KP, Dosemeci M, et al. Occupational and leisure-time physical activity and risk of colon cancer by subsite. *J Occup Environ Med* 2006 ; 48 : 236-43.

Zheng W, Shu XO, McLaughlin JK, et al. Occupational physical activity and the incidence of cancer of the breast, corpus uteri, and ovary in Shanghai. *Cancer* 1993 ; 71 : 3620-4.

ESPÉRANCE DE VIE

Alevizos A, Lentzas J, Kokkoris S, et al. Physical activity and stroke risk. *Int J Clin Pract* 2005 ; 59 : 922-30.

Alfano CM, Klesges RC, Murray DM, et al. Physical activity in relation to all-site and lung cancer incidence and mortality in current and former smokers. *Cancer Epidemiol Biomarkers Prev* 2004 ; 13 : 2233-41.

Andersen LB, Schnohr P, Schroll M, Hein HO. All-Cause Mortality Associated With Physical Activity During Leisure Time, Work, Sports, and Cycling to Work. *Arch Intern Med*. 2000 ; 160 : 1621-8.

Barengo NC, Hu G, Lakka TA, et al. Low physical activity as a predictor for total and cardiovascular disease mortality in middle-aged men and women in Finland. *Eur Heart J* 2004 ; 25 : 2204-11.

Crespo CJ, Palmieri MR, Perdomo RP, et al. The relationship of physical activity and body weight with all-cause mortality : results from the Puerto Rico Heart Health Program. *Ann Epidemiol* 2002 ; 12 : 543-52.

Davis MA, Neuhaus JM, Moritz DJ, et al. Health behaviors and survival among middle-aged and older men and women in the NHANES I Epidemiologic Follow-up Study. *Prev Med* 1994 ; 23 : 369-76.

Eaton CB, Medalie JH, Flocke SA, et al. Self-reported physical activity predicts long-term coronary heart disease and all-cause mortalities. Twenty-one year follow-up of the Israeli ischemic heart disease study. *Arch Fam Med* 1995 ; 4 : 323-9.

Ellekjær EF, Wyller TB, Sverre JM, Holmen J. Lifestyle factors and risk of cerebral infarction. *Stroke* 1992 ; 23 : 829-34.

Ellekjær H, Holmen J, Ellekjær E, Vatten L. Physical activity and stroke mortality in women. Ten-year follow-up of the Nord-Trøndelag health survey, 1984-1986. *Stroke* 2000 ; 31 : 14-8.

France Portrait social (éditions 2005-2006). Paris : Insee éditions, 2005.

Fujita K, Takahashi H, Miura C, et al. Walking and mortality in Japan : the Miyagi Cohort Study. *J Epidemiol* 2004 ; 14 : S26-32.

Gillum RF, Mussolino ME, Ingram DD. Physical activity and stroke incidence in women and men : the NHANES I Epidemiologic Follow-up Study. *Am J Epidemiol* 1996 ; 143 : 860-9.

Giovannucci EL, Liu Y, Leitzmann MF, et al. A prospective study of physical activity and incident and fatal prostate cancer. *Arch Intern Med* 2005 ; 165 : 1005-10

Gregg EW, Cauley JA, Stone K, et al. Relationship of changes in physical activity and mortality among older women. *JAMA* 2003 ; 289 : 2379-86.

Haapanen-Niemi N, Miilunpalo S, Pasanen M, et al. Body mass index, physical activity and low level of physical fitness as determinants of all-cause and cardiovascular disease mortality-16 y follow-up of middle-aged and elderly men and women. *Int J Obes Relat Metab Disord* 2000 ; 24 : 1465-74.

Haut Comité de la Santé Publique. *La santé en France 2002*. Paris : Ministère de l'Emploi et de la Solidarité, janvier 2002.

Holmes MD, Chen WY, Feskanich D, et al. Physical activity and survival after breast cancer diagnosis. *JAMA* 2005 ; 293 : 2479-86.

Hu G, Jousilahti P, Barengo NC, et al. Physical Activity, Cardiovascular Risk Factors, and Mortality Among Finnish Adults With Diabetes. *Diabetes Care* 2005 ; 28 : 799-805.

Hu G, Tuomilehto J, Silventoinen K, et al. The effects of physical activity and body mass index on cardiovascular, cancer and all-cause mortality among 47 212 middle-aged Finnish men and women. *Int J Obes Relat Metab Disord* 2005 ; 29 : 894-902.

Janssen I, Jolliffe CJ. Influence of physical activity on mortality in elderly with coronary artery disease. *Med Sci Sports Exerc* 2006 ; 38 : 418-7.

Jonker JT, De Laet C, Franco OH, et al. Physical activity and life expectancy with and without diabetes : life table analysis of the Framingham Heart Study. *Diabetes Care* 2006 ; 29 : 38-43.

Kampert JB, Blair SN, Barlow CE, et al. Physical activity, physical fitness, and all-cause and cancer mortality : a prospective study of men and women. *Ann Epidemiol* 1996 ; 6 : 452-7.

Kaplan GA, Strawbridge WJ, Cohen RD, et al. Natural history of leisure-time physical activity and its correlates : associations with mortality from all causes and cardiovascular disease over 28 years. *Am J Epidemiol* 1996 ; 144 : 793-7.

Kiely DK, Wolf AW, Cupples LA. Physical activity and stroke risk : the Framingham Study. *Am J Epidemiol* 1994 ; 140 : 608-20.

Kilander L, Berglund L, Boberg M, et al. Education, lifestyle factors and mortality from cardiovascular disease and cancer. A 25-year follow-up of Swedish 50-year-old men. *Int J Epidemiol* 2001 ; 30 : 1119-26.

Kristal-Boneh E, Harari G, Melamed S, Froom P. Association of Physical Activity at Work With Mortality in Israeli Industrial Employees : The CORDIS Study. *J Occup Env Med* 2000 ; 42 : 127-35.

Kujala UM, Kaprio J, Sarna S, Koskenvuo M. Relationship of leisure-time physical activity and mortality : the Finnish twin cohort. *JAMA* 1998 ; 279 : 440-4.

Kushi LH, Fee RM, Folsom AR, et al. Physical activity and mortality in postmenopausal women. *JAMA* 1997 ; 277 : 1287-92.

Lam TH, Ho SY, Hedley AJ, et al. Leisure time physical activity and mortality in Hong Kong : case-control study of all adult deaths in 1998. *Ann Epidemiol* 2004 ; 14 : 391-8.

Landi F, Cesari M, Onder G, et al. Physical activity and mortality in frail, community-living elderly patients. *J Gerontol A Biol Sci Med Sci* 2004 ; 59 : 833-7.

Lee IM, Paffenbarger RS. Physical activity and stroke incidence : the Harvard Alumni Health Study. *Stroke* 1998 ; 29 : 2049 -54.

Lee IM, Sesso HD, Oguma Y, Paffenbarger RS Jr. Physical activity, body weight, and pancreatic cancer mortality. *Br J Cancer* 2003 ; 88 : 679-83.

Lee Y. The predictive value of self assessed general, physical, and mental health on functional decline and mortality in older adults. *J Epidemiol Community Health* 2000 ; 54 : 123-9.

Manini TM, Everhart JE, Patel KV, et al. Daily activity energy expenditure and mortality among older adults. *J Am Med Assoc* 2006 ; 296 : 171-9.

Mensink GB, Deketh M, Mul MD, et al. Physical activity and its association with cardiovascular risk factors and mortality. *Epidemiology* 1996 ; 7 : 391-7.

Myers J, Kaykha A, George S, et al. Fitness versus physical activity patterns in predicting mortality in men. *Am J Med* 2004 ; 117 : 912-8.

O'Connor GT, Buring JE, Yusuf S, et al. An overview of randomized trials of rehabilitation with exercise after myocardial infarction. *Circulation* 1989 ; 80 : 234 -44.

Paffenbarger RS Jr, Hyde RT, Wing AL. Physical activity and incidence of cancer in diverse populations : a preliminary report. *Am J Clin Nutr* 1987 ; 45 : 312-7.

Paffenbarger RS, Hyde RT, Wing AL, et al. Physical activity, all-cause mortality, and longevity of college alumni. *N Engl J Med* 1986 ; 314 : 605-13.

Pekkanen J, Marti B, Nissinen A, Tuomilehto J, Punsar S, Karvonen MJ. Reduction of premature mortality by high physical activity : a 20-year follow-up of middleaged Finnish men. *Lancet* 1987 ; 1 : 1473-7.

Polednak AP. College athletes, body size, and cancer mortality. *Cancer* 1976 ; 38 : 382-7.

Richardson CR, Kriska AM, Lanz P, Hayward RA. Physical Activity and Mortality across Cardiovascular Disease Risk Groups. *Med Sci Sports Exerc* 2004 ; 36 : 1923-9.

Richet-Mastain L. Bilan démographique 2005. *Insee Première* 2006 ; 1059 : 1-4.

Rohan TE, Fu W, Hiller JE. Physical activity and survival from breast cancer. *Eur J Cancer Prev* 1995 ; 4 : 419-24.

Rosengren A, Wilhelmsen L. Physical activity protects against coronary death and deaths from all causes in middle-aged men. Evidence from a 20-year follow-up of the primary prevention study in Goteborg. *Ann Epidemiol* 1997 ; 7 : 69-75.

Sacco RL, Gan R, Boden-Albala B, et al. Leisure-time physical activity and ischemic stroke risk : the Northern Manhattan Stroke Study. *Stroke* 1998 ; 29 : 380-7.

Sarna S, Kaprio J, Kujala UM, Koskenvuo M. Health status of former elite athletes. The Finnish experience. *Aging* 1997 ; 9 : 35-41.

Sarna S, Sahi T, Koskenvuo M, Kaprio J. Increased life expectancy of world class male athletes. *Med Sci Sports Exerc* 1993 ; 25 : 237-44.

Schnohr P, Scharling H, Jensen JS. Changes in Leisure-time Physical Activity and Risk of Death : An Observational Study of 7,000 Men and Women. *Am J Epidemiol* 2003 ; 158 : 639-44.

Sherman SE, D'Agostino RB, Silbershatz H, Kannel WB. Comparison of past versus recent physical activity in the prevention of premature death and coronary artery disease. *Am Heart J* 1999 ; 138 : 900-7.

Stessman J, Maaravi Y, Hammerman-Rozenberg R, et al. The effects of physical activity on mortality in the Jerusalem 70-Year-Olds Longitudinal Study. *J Am Geriatr Soc* 2000 ; 48 : 499-504.

Sundquist K, Qvist J, Sundquist J, Johansson SE. Frequent and occasional physical activity in the elderly : a 12-year follow-up study of mortality. *Am J Prev Med* 2004 ; 27 : 22-7.

Thune I, Lund E. The influence of physical activity on lung-cancer risk : a prospective study on 81516 men and women. *Int J Cancer* 1997 ; 70 : 57-62.

Trichopoulos A, Psaltopoulou T, Orfanos P, Trichopoulos D. Diet and physical activity in relation to overall mortality amongst adult diabetics in a general population cohort. *J Intern Med* 2006 ; 259 : 583-91.

Villeneuve PJ, Morrison HI, Craig CL, et al. Physical activity, physical fitness, and risk of dying. *Epidemiol* 1998 ; 9 : 626-31 ;

Wannamethee SG, Shaper AG, Walker M. Physical Activity and Mortality in Older Men With Diagnosed Coronary Heart Disease. *Circulation* 2000 ; 102 : 1358-63.

Weller I, Corey P. The impact of excluding non-leisure energy expenditure on the relation between physical activity and mortality in women. *Epidemiology* 1998 ; 9 : 632-5.

WHO. *Health Promotion Glossary*. WHO/HPR/HEP/98.1. Geneva : WHO, 1999

Williams PT. Health effects resulting from exercise versus those from body fat loss. *Med Sci Sports Exerc* 2001 ; 33 : S611-21.

Wyshak G. Behavioral practices and mortality in women former college athletes and nonathletes. *Health Care Women Int* 2003 ; 24 : 808-21.

QUALITÉ DE VIE

Baatile J, Langbein WE, Weaver F, et al. Effect of exercise on perceived quality of life of individuals with Parkinson's disease. *J Rehabil Res Develop* 2000 ; 37 : 529-34.

Brooks R, Rabin RE, de Charro Fth. *The measurement and valuation of health status using EQ-5D : a European perspective*. Kluwer Academic Publishers : 2003.

Brown DW, Balluz LS, Heath GW, et al. Associations between recommended levels of physical activity and health-related quality of life. *Prev Med* 2003 ; 37 : 520-8.

Brown DW, Brown DR, Heath GW, et al. Associations between physical activity dose and health-related quality of life. *Med Sci Sports Exerc* 2004 ; 36 : 890-6.

Bruchon-Schweitzer M. *Psychologie de la santé. Modèles, concepts et méthodes*. Paris : Dunod, 2002 : 43-84.

Bui-Xuan G, Brunet F, Dejean O. Activité physique, santé et qualité de vie : Enquête sur l'activité physique, la santé et la qualité de vie des personnes déficientes intellectuelles de CAT. *Handicap* 2000 ; 85 : 47-67.

Calmels P. Douleur chronique, activité physique et qualité de vie. *Sci Sports* 1998 ; 13 : 10-6.

Campbell A, Converse C, Rodgers W. *The quality of American life*. New York : Russel Sage. 1976.

Chen X, Sekine M, Hamanishi S, et al. Lifestyles and health-related quality of life in Japanese school children : a cross-sectional study. *Prev Med* 2005 ; 40 : 668-78.

Chen X, Sekine M, Hamanishi S, Yamagami T, Kagamimori S. Associations of lifestyle factors with quality of life (QOL) in Japanese children : a 3-year follow-up of the Toyama Birth Cohort Study. *Child Care Health Dev* 2005 ; 31 : 433-9.

Ferrari M, Vangelista A, Vedovi E, et al. Minimally supervised home rehabilitation improves exercise capacity and health status in patients with COPD. *Am J Phys Med Rehabil* 2004 ; 83 : 337-43.

Gérin P, Dazord A, Cialdella P, et al. Le questionnaire profil de la qualité de vie subjective. *Thérapie* 1991 ; 46 : 131-8.

Glasgow RE, Ruggiero L, Eakin EG, et al. Quality of life and associated characteristics in a large national sample of adults with diabetes. *Diabetes Care* 1997 ; 20 : 562-7.

Greenlund KJ, Giles WH, Keenan NL, et al. Physician advice, patient actions, and health-related quality of life in secondary prevention of stroke through diet and exercise. *Stroke* 2002 ; 33 : 565-70

Hotlon R, Colberg SR, Nunnold T, et al. The Effect of an Aerobic Exercise Training Program on Quality of Life in Type 2 Diabetes. *Diab Educator* 2003 ; 29 : 837-46.

- Hulens M, Vansant G, Claessens AL, et al. Health-related quality of life in physically active and sedentary obese women. *Am J Hum Biol* 2002 ; 14 : 777-85.
- Ishizawa Y, Chogahara W. An empirical study on the determinants of life satisfaction among elderly Japanese. *2000 Pre-Olympic Congress. Sports medicine and physical education. International Congress on Sport Science. 7-13 september, Brisbane, Australia 2000.*
- Jolliffe JA, Rees K, Taylor RS, et al. Exercise-based rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* 2001 ; 1 : CD001800.
- Kaplan RM, Bush JW. Health-related quality of life measurement for evaluation research and policy analysis. *Health Psychol* 1982 ; 1 : 61-80.
- Khanna S, Pal H, Pandey RM, Handa R. The relationship between disease, activity and quality of life in systemic lupus erythematosus. *Rheumatology* 2004 ; 43 : 1536-40.
- Kovac D. Quality of the environment and quality of life. In : Georgas J, Manthouli M, Besevegis E, Kokkeki A. *Contemporary psychology in Europe. Theory, research and applications.* Seattle : Hogrefe et Huber, 1996.
- Launois R. La qualité de vie : finalités et méthodes. In : Moatti JP. *Recherche clinique et qualité de vie.* Paris : Flammarion, 1996 : 61-8.
- Lustky KB, Widman L, Paschane AAE, Oslon KC. Physical activity and quality of life : assessing the influence of activity frequency, intensity, volume, and motives. *Behavioral Med* 2004 ; 22 september.
- McAuley E, Rudolph D. Physical activity, aging, and psychological well-being. *J Aging Phys Activity* 1995 ; 3 : 67-96.
- Millat B. Mesure de la qualité de vie comme critère de jugement en chirurgie générale et digestive. In : Moatti JP. *Recherche clinique et qualité de vie.* Paris : Flammarion, 1996 : 121-8.
- Morimoto T, Oguma Y, Yamazaki S, et al. Gender differences in effects of physical activity on quality of life and resource utilization. *Qual Life Res* 2006 ; 15 : 537-46.
- Nelson EU. The COOP Charts. In : McDowell I, Newell C, editors. *Measuring health : a guide to rating scales and questionnaire.* Second ed. New York Univ Press ; 1996 : 410-6
- Oudry A, Hermand D. Qualité de vie urbaine : conceptualisation et évaluation. In : Bonardi C, Girandola F, Roussiau N, Soubiale N. *Psychologie sociale appliquée. Environnement, santé, qualité de vie.* Paris : In press Editions, 2002 : 201-16.
- Painter P, Carlson L, Carey S, et al. Physical functioning and health-related quality-of-life changes with exercise training in hemodialysis patients. *Am J Kidney Dis* 2000 ; 35 : 482-92.
- Prior JC, Barr SI, Chow LR, Faulkner RA. Prevention and management of osteoporosis : consensus statements from the Scientific Advisory Board of the Osteoporosis Society of Canada. Physical activity as therapy for osteoporosis. *Can Med Assoc J* 1996 ; 155 : 940-44.
- Rees K, Taylor RS, Singh S, et al. Exercise based rehabilitation for heart failure. *The Cochrane Database Syst Rev* 2004 ; (3) : CD003331.
- Rizzo L, Spitz E. Qualité de vie, santé et maladie. In : Fischer GN. *Traité de psychologie de la santé.* Paris : Dunod, 2002 : 283-99.
- Rorth M, Andersen C, Quist M, et al. Health benefits of a multidimensional exercise program for cancer patients undergoing chemotherapy. *ASCO Meeting Abstracts* 2005 ; 23 : 8010.
- Steptoe A, Butler N. Sports participation and emotional wellbeing in adolescents. *Lancet* 1996 ; 347 : 1789-92.
- The EuroQoL Group. EuroQoL-a new facility for the measurement of health-related quality of life. *Health Policy* 1990 ; 16 : 199-208.
- Thorsen L, Skovlund E, Strømme SB, et al. Effectiveness of Physical Activity on Cardiorespiratory Fitness and Health-Related Quality of Life in Young and Middle-Aged Cancer Patients Shortly After Chemotherapy. *J Clin Oncol* 2005 ; 23 : 2378-88.
- Villaverde-Gutierrez C, Araujo E, Cruz F, et al. Quality of life of rural menopausal women in response to a customized exercise programme. *J Adv Nurs* 2006 ; 54 : 11-9.
- Vlek C, Skolnik M, Gatersleben B. Sustainable development and quality of life : expected effects of prospective changes in economic and environmental conditions. *Z Exp Psychol* 1998 ; 45 : 319-33.
- Vuillemin A, Boini S, Bertrais S, et al. Leisure time physical activity and health-related quality of life. *Prev Med* 2005 ; 41 : 562-9.
- Ware JE, Gandek B. Overview of the SF-36 Health Survey and the International Quality of Life Assessment (IQOLA) project. *J Clin Epidemiol* 1998 ; 51 : 903-12.
- Wendel-Vos GC, Schuit AJ, Tijhuis MA, Kromhout D. Leisure time physical activity and health-related quality of life : cross-sectional and longitudinal associations. *Qual Life Res* 2004 ; 13 : 667-77.
- World Health Organization. *WHOQoL Study Protocol.* WHO, MNH7PSF/93.9, 1993.



Dangers de l'activité physique, éléments pour leur maîtrise

MORTALITÉ

- Ades PA. Preventing sudden death : cardiovascular screening of young athletes. *Phys Sports Med* 1992 ; 20 : 75-89.
- Baker SP, Li G, Fowler C, Dannenberg AL. *Injuries to bicyclists : a national perspective*. Monograph prepared for the Snell Memorial Foundation. Baltimore (MD) : Johns Hopkins University Injury Prevention Center, 1993.
- Basilico FC. Cardiovascular diseases in athletes. *Am J Sports Med* 1999 ; 27 : 108-21.
- Carré F. Bénéfices et risques de la pratique d'une activité physique. *Ann Cardiol Angéiol* 2002 ; 51 : 351-6.
- Corrado D, Basso C, Rizzoli G, Schiavon M, Thiene G. Does sports activity enhance the risk of sudden death in adolescents and young adults ? *J Am Coll Cardiol* 2003 ; 42 : 1959-63
- Dupuis JM, Tabib A, Reix P, et al. Mort subite inopinée d'origine cardiaque entre 6 et 18 ans. Données anatomopathologiques. Rôle du sport ? Prévention ? *Arch Ped* 2005 ; 12 : 1204-8.
- Ekman R, Welander G, Svanstrom L, et al. Bicycle-related injuries among the elderly – a new epidemic ? *Public Health* 2001 ; 115 : 38-43
- Fornes P, Lecomte D. Mort subite et activité physique et sportive. *Rev Prat* 2001 ; 51 : S31-5.
- Gabbe BJ, Finch CF, Cameron PA, Williamson OD. Incidence of serious injury and death during sport and recreation activities in Victoria, Australia. *Br J Sports Med* 2005 ; 39 : 573-7
- Gardner JW, Gutmann FD, Potter RN, Kark JA. Nontraumatic exercise-related deaths in the U.S. military, 1996-1999. *Mil Med* 2002 ; 167 : 964-70.
- INED. *Population & Sociétés* 2005 ; 414 : 3.
- Leski M. Sudden cardiac death in athletes. *South Med J* 2004 ; 97 : 861-2.
- Lindholm P, Steensberg J. Epidemiology of unintentional drowning and near-drowning in Denmark in 1995. *Inj Prev* 2000 ; 6 : 29-31.
- Maron BJ, Gohman TE, Aeppli D. Prevalence of sudden cardiac death during competitive sports activities in Minnesota high school athletes. *J Am Coll Cardiol* 1998 ; 32 : 1881-4.
- Maron BJ, Poliac LC, Roberts WO. Risk of sudden cardiac death associated with marathon running. *J Am Coll Cardiol* 1996 ; 28 : 428-31.
- Maron BJ. Sudden death in young athletes. *N Eng J Med* 2003 ; 349 : 1064-75.
- Missoni E, Kern J. Fatality risk factors for bicyclists in Croatia. *Croat Med J* 2003 ; 44 : 610-3.
- Osberg JS, Stiles SC, Asare OK. Bicycle safety behavior in Paris and Boston. *Accid Anal Prev* 1998 ; 30 : 679-87.
- Quigley F. A survey of the causes of sudden death in sport in the Republic of Ireland. *Br J Sports Med* 2000 ; 34 : 258-61.
- Ragosta M, Crabtree J, Sturner WQ, Thompson PD. Death during recreational exercise in the State of Rhode Island. *Med Sci Sports Exerc.* 1984 ; 16 : 339-42.
- Thompson MJ, Rivara FP. Bicycle-related injuries. *Am Fam Physician* 2001 ; 63 : 2007-14.

MORBIDITÉ

- Belechri M, Petridou E, Kedikoglou S, et al. Sports injuries among children in six European union countries. *Eur J Epidemiol* 2001 ; 17 : 1005-12.
- Cassell EP, Finch CF, Stathakis VZ. Epidemiology of medically treated sport and active recreation injuries in the Latrobe Valley, Victoria, Australia. *Br J Sports Med* 2003 ; 37 : 405-9.
- Centers for Disease Control and Prevention (CDC). Nonfatal sports- and recreation-related injuries treated in emergency departments-United States, July 2000-June 2001. *MMWR Morb Mortal Wkly Rep* 2002 ; 51 : 736-40.

Chapuis R, Schléret J.-M. Rapport annuel 2003. Paris : Observatoire national de la sécurité des établissements scolaires et d'enseignement supérieur, 2004.

Conn JM, Annett JL, Gilchrist J. Sports and recreation related injury episodes in the US population, 1997-99. *Inj Prev* 2003 ; 9 : 117-23.

Daigle J.-M. *Hospitalisations pour traumatismes d'origine récréative et sportive au Québec de 1994 à 2002*. Institut national de santé public du Québec, novembre 2004.

Dempsey RL, Layde PM, Laud PW, et al. Incidence of sports and recreation related injuries resulting in hospitalization in Wisconsin in 2000. *Inj Prev* 2005 ; 11 : 91-6.

Gary F. Les accidents de sport chez les jeunes de 10 à 24 ans. *Point Stat (Cnamts)* 1999 ; 14 : 1-4.

Gerson LW, Stevens JA. Recreational injuries among older Americans, 2001. *Inj Prev* 2004 ; 10 : 134-8.

Kelly KD, Lissel HL, Rowe BH, et al. Sport and recreation-related head injuries treated in the emergency department. *Clin J Sport Med* 2001 ; 11 : 77-81.

Mackie IJ. Patterns of drowning in Australia, 1992-1997. *Med J Aust* 1999 ; 171 : 587-90.

Michaud PA, Renaud A, Narring F. Sports activities related to injuries ? A survey among 9-19 year olds in Switzerland. *Inj Prev* 2001 ; 7 : 41-5.

Parkkari J, Kannus P, Natri A, et al. Active living and injury risk. *Int J Sports Med* 2004 ; 25 : 209-16.

Robinson DL. Safety in numbers in Australia : more walkers and bicyclists, safer walking and bicycling. *Health Promot J Austr* 2005 ; 16 : 47-51.

Schneider S, Seither B, Tonges S, Schmitt H. Sports injuries : population based representative data on incidence, diagnosis, sequelae, and high risk groups. *Br J Sports Med* 2006 ; 40 : 334-9.

Spinks AB, McClure RJ, Bain C, Macpherson AK. Quantifying the association between physical activity and injury in primary school-aged children. *Pediatrics* 2006 ; 118 : e43-50.

Stevenson MR, Hamer P, Finch CF, et al. Sport, age, and sex specific incidence of sports injuries in Western Australia. *Br J Sports Med* 2000 ; 34 : 188-94.

Thélot B, Ricard C. *Résultats de l'Enquête permanente sur les accidents de la vie courante, années 2002-2003*. Réseau Epac. Institut de veille sanitaire, octobre 2005.

ELÉMENTS DE MAÎTRISE DES DANGERS

American College of Sports Medicine. Position stand on exercise and fluid replacement. *Med Sci Sports Exerc* 1996 ; 28 : i-vii.

Andersen JC. Stretching before and after exercise : effect on muscle soreness and injury risk. *J Athl Train* 2005 ; 40 : 218-20.

Attewell RG, Glase K, McFadden M. Bicycle helmet efficacy : a meta-analysis. *Accid Anal Prev* 2001 ; 33 : 345-52.

Bernheim A. Höhe und Herzkrankheiten. *Schweiz Rundsch Med Prax* 2005 ; 45 : 1760-4.

Bishop D. Warm up II : performance changes following active warm up and how to structure the warm up. *Sports Med* 2003 ; 33 : 483-98.

Bourdessol H, Gautier A, Guilbert P et al. Pratique du roller et port du casque. *Bull Epidemiol Hebdo* 2001 ; 13 : 1-4.

Campbell ME, Li Q, Gingrich SE, et al. Should people be physically active outdoors on smog alert days ? *Can J Public Health* 2005 ; 96 : 24-8.

Carek PJ, Mainous A. Preparticipation physical examination for athletics : a systematic review of current recommendations. *Br J Med* 2003 ; 327 : 170-3.

Carter R, Watenpaugh DE, Smith ML. Gender differences in cardiovascular regulation during recovery from exercise. *J Appl Physiol* 2001 ; 91 : 1902-7.

Casa DJ, Armstrong LE, Hillman SK, et al. National Athletic Trainers' Association Position Statement : Fluid replacement for athletes. *J Athl Train* 2000 ; 35 : 212-24.

Chertok M, Voukelatos A, Sheppard V, Rissel C. Comparison of air pollution exposure for five commuting modes in Sydney – car, train, bus, bicycle and walking. *Health Promot J Aust* 2004 ; 15 : 63-7.

Coris EE, Ramirez AM, Van Durme DJ. Heat illness in athletes : the dangerous combination of heat, humidity and exercise. *Sports Med* 2004 ; 34 : 9-16.

Courneya KS, Mackey JR, Jones LW. Coping with cancer. Can exercise help ? *Phys Sportsmed* 2000 ; 28 (5) : on line.

Daigle CC, Chalupa DC, Gibb FR, et al. Ultrafine particle deposition in humans during rest and exercise. *Inhal Toxicol* 2003 ; 15 : 539-52.

Dib MY, Smith J, Bernhardt KA, et al. Effect of environmental temperature on shock absorption properties of running shoes. *Clin J Sport Med* 2005 ; 15 : 172-6.

Donaldson GC, Keatinge WR. Early increases in ischaemic heart disease mortality dissociated from and later changes associated with respiratory mortality after cold weather in south east England. *J Epidemiol Community Health* 1997 ; 51 : 643-8.

Donaldson GC, Rintamaki H, Nayha S. Outdoor clothing : its relationship to geography, climate, behaviour and cold-related mortality in Europe. *Int J Biometeorol* 2001 ; 45 : 45-51.

Ehrlich PF, Longhi J, Vaughan R, Rockwell S. Correlation between parental perception and actual childhood patterns of bicycle helmet use and riding practices : implications for designing injury prevention strategies. *J Pediatr Surg* 2001 ; 36 : 763-6.

Evans RK, Knight KL, Draper DO, Parcell AC. Effects of warm-up before eccentric exercise on indirect markers of muscle damage. *Med Sci Sports Exerc* 2002 ; 34 : 1892-9.

Faigenbaum AD, Bellucci M, Bernieri A, et al. Acute effects of different warm-up protocols on fitness performance in children. *J Strength Cond Res* 2005 ; 19 : 376-81.

Forjuoh SN, Schuchmann JA, Fiesinger T, Mason S. Parent-child concordance on reported barriers to helmet use by children. *Med Sci Monit* 2003 ; 9 : CR436-41.

Gibbons RJ, Balady GJ, Bricker JT, et al. ACC/AHA 2002 guideline update for exercise testing : summary article : a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1997 Exercise Testing Guidelines). *Circulation* 2002 ; 106 : 1883-92

Hart L. Effect of stretching on sport injury risk : a review. *Clin J Sport Med* 2005 ; 15 : 113.

Herbert RD, Gabriel M. Effects of stretching before and after exercising on muscle soreness and risk of injury : systematic review. *Br Med J* 2002 ; 325 : 468.

Hildebrandt W, Schutze H, Stegemann J. Cardiovascular limitations of active recovery from strenuous exercise. *Eur J Appl Physiol Occup Physiol* 1992 ; 64 : 250-7.

Hue O, Voltaire B, Galy O, et al. Effects of 8 days acclimation on biological and performance response in a tropical climate. *J Sports Med Phys Fitness* 2004 ; 44 : 30-7.

Jagodzinski T, DeMuri GP. Horse-related injuries in children : a review. *WMJ* 2005 ; 104 : 50-4.

Kennedy A. The pattern of injury in fatal pedal cycle accidents and the possible benefits of cycle helmets. *Br J Sports Med* 1996 ; 30 : 130-3.

Koenig JQ, Pierson WE, Horike M, et al. Effects of SO₂ plus NaCl aerosol combined with moderate exercise on pulmonary function in asthmatic adolescents. *Environ Res* 1981 ; 25 : 340-8.

Maeda T, Sugawara A, Fukushima T, et al. Effects of lifestyle, body composition, and physical fitness on cold tolerance in humans. *J Physiol Anthropol Appl Human Sci* 2005 ; 24 : 439-43.

Mandengue SH, Seck D, Bishop D, et al. Are athletes able to self-select their optimal warm up ? *J Sci Med Sport* 2005 ; 8 : 26-34.

Parkkari J, Kujala UM, Kannus P. Is it possible to prevent sports injuries ? Review of controlled clinical trials and recommendations for future work. *Sports Med* 2001 ; 31 : 985-95.

Pekkanen J, Peters A, Hoek G, et al. Particulate air pollution and risk of ST-segment depression during repeated submaximal exercise tests among subjects with coronary heart disease : the exposure and risk assessment for fine and ultrafine particles in ambient air (ULTRA) study. *Circulation* 2002 ; 106 : 933-8.

Ponchia A, Biasin R, Tempesta T, et al. Cardiovascular risk during physical activity in the mountains. *J Cardiovasc Med* 2006 ; 7 : 129-35.

Pope CA 3rd, Burnett RT, Thurston GD, et al. Cardiovascular mortality and long-term exposure to particulate air pollution : epidemiological evidence of general pathophysiological pathways of disease. *Circulation* 2004 ; 109 : 71-7.

Reisman S, Walsh LD, Proske U. Warm-up stretches reduce sensations of stiffness and soreness after eccentric exercise. *Med Sci Sports Exerc* 2005 ; 37 : 929-36.

Ruidavets JB, Cassadou S, Cournot M, et al. Increased resting heart rate with pollutants in a population based study. *J Epidemiol Community Health* 2005 ; 59 : 685-93.

- Saat M, Sirisinghe RG, Singh R, Tochihara Y. Effects of short-term exercise in the heat on thermoregulation, blood parameters, sweat secretion and sweat composition of tropic-dwelling subjects. *J Physiol Anthropol Appl Human Sci* 2005 ; 24 : 541-9.
- Stamford BA, Weltman A, Moffatt R, Sady S. Exercise recovery above and below anaerobic threshold following maximal work. *J Appl Physiol* 1981 ; 51 : 840-4.
- Stocks JM, Taylor NA, Tipton MJ, Greenleaf JE. Human physiological responses to cold exposure. *Aviat Space Environ Med* 2004 ; 75 : 444-57.
- Sulheim S, Holme I, Ekeland A, Bahr R. Helmet use and risk of head injuries in alpine skiers and snowboarders. *JAMA* 2006 ; 295 : 919-24.
- Taylor D, Fergusson M. The comparative pollution exposure of road users – a summary. *World Transp Pol Pract* 1998 ; 4 : 22-6.
- Taylor DM, Bennett D, Carter M, et al. Perceptions of surfboard riders regarding the need for protective headgear. *Wilderness Environ Med.* 2005 ; 16 : 75-80.
- Thacker SB, Gilchrist J, Stroup DF, Kimsey CD Jr. The impact of stretching on sports injury risk : a systematic review of the literature. *Med Sci Sports Exerc* 2004 ; 36 : 371-8.
- Threshold limit values and biological exposure indices, 2001.* Cincinnati : American Conference of Governmental Industrial Hygienists, 2001.
- Von Duvillard SP, Braun WA, Markofski M, et al. Fluids and hydration in prolonged endurance performance. *Nutrition* 2004 ; 20 : 651-6.
- Weerapong P, Hume PA, Kolt GS. The mechanisms of massage and effects on performance, muscle recovery and injury prevention. *Sports Med* 2005 ; 35 : 235-56.
- West JB. The physiologic basis of high-altitude diseases. *Ann Intern Med* 2004 ; 141 : 789-800
- Wigernaes I, Hostmark AT, Stromme SB, et al. Active recovery and post-exercise white blood cell count, free fatty acids, and hormones in endurance athletes. *Eur J Appl Physiol* 2001 ; 84 : 358-66.
- Wingfield K, Matheson GO, Meeuwisse WH. Preparticipation evaluation : an evidence-based review. *Clin J Sport Med* 2004 ; 14 : 109-22.
- Withnall R, Eastaugh J, Freemantle N. Do shock absorbing insoles in recruits undertaking high levels of physical activity reduce lower limb injury ? A randomized controlled trial. *J R Soc Med* 2006 ; 99 : 32-7.
- Witvrouw E, Mahieu N, Danneels L, McNair P. Stretching and injury prevention : an obscure relationship. *Sports Med* 2004 ; 34 : 443-9.
- Zainuddin Z, Newton M, Sacco P, Nosaka K. Effects of massage on delayed-onset muscle soreness, swelling, and recovery of muscle function. *J Athl Train* 2005 ; 40 : 174-80.



Engagement durable dans une activité physique

DÉTERMINANTS INDIVIDUELS

- Abramson S, Stein J, Schaufele M, et al. Personal exercise habits and counseling practices of primary care physicians : a national survey. *Clin J Sport Med* 2000 ; 10 : 40-8.
- Ajzen I. *Attitudes, personality and behaviour.* Milton Keynes : Open university press, 1991
- Al-Ali N, Haddad LG. The effect of the health belief model in explaining exercise participation among Jordanian myocardial infarction patients. *J Transcult Nurs* 2004 ; 15 : 114-21.
- American College of Sports Medicine RM. *ACSM's guidelines for exercise testing and prescription*, sixth ed. Philadelphia (PA) Lippincott, Williams, Wilkins : 2000
- Anderson CB. When more is better : number of motives and reasons for quitting as correlates of physical activity in women. *Health Education Res* 2003 ; 18 : 525-37.

Baer N. MDs seen as key players in move to encourage active living in Canada. *CMAJ* 1997 ; 156 : 1605-8.

Baerschi B. L'homme et son espèce : l'espèce humaine est-elle menacée par les progrès des biotechnologies ? in : Huber G, Byk C. *La bioéthique au pluriel*. Paris : John Libbey, 1996 : 17-30.

Balde A, Figueras J, Hawking DA, et al. Physician advice to the elderly about physical activity. *JAPA* 2003 ; 11 : 90-7.

Bandura A. *Social foundations of thought and action : a social-cognitive theory*. Englewood Cliffs : Prentice-Hall, 1986.

Bandura A. *Social learning theory*. Englewood Cliffs : Prentice-Hall, 1977.

Bauman AE, Russell SJ, Furber SE, Dobson AJ. The epidemiology of dog walking : an unmet need for human and canine health. *Med J Aust* 2001 ; 175 : 632-4.

Bauman AE, Sallis JF, Dzawaltowski DA, Owen N. Toward a better understanding of the influences on physical activity. The role of determinants, correlates, causal variables, mediators, moderators, and confounders. *Am J Prev Med* 2002 ; 23 : 5-14.

Biddle S, Goudas M. Analysis of children's physical activity and its association with adult encouragement and social cognitive variables. *J School Health* 1996 ; 66 : 75-8.

Bois JE, Sarrazin PG, Brustad RJ, et al. Elementary schoolchildren's perceived competence and physical activity involvement : the influence of parents' role modelling behaviours and perceptions of their child's competence. *Psychol Sport Exerc* 2005 ; 6 : 381-97.

Booth ML, Owen N, Bauman A, et al. Social-cognitive and perceived environment influences associated with physical activity in older Australians. *Prev Med* 2000 ; 31 : 15-22.

Bortons C, Björkelund C, Bulc M, et al. Prevention and health promotion in clinical practice : the views of general practitioners in Europe. *Prev Med* 2005 ; 40 : 595-601.

Brown SA, Huber D, Bergman A. A perceived Benefits and Barriers Scale for strenuous physical activity in college students. *Am J Health Promot* 2006 ; 21 : 137-40.

Brown SA. Measuring perceived benefits and perceived barriers for physical activity. *Am J Health Behav* 2005 ; 29 : 107-16.

Brown SG, Rhodes RE. Relationships among dog ownership and leisure-time walking in Western Canadian adults. *Am J Prev Med* 2006 ; 30 : 131-6.

Brownson RC, Baker EA, Housemann RA, et al. Environmental and Policy Determinants of Physical Activity in the United States. *Am J Public Health* 2001 ; 91 : 1995-2003.

Brunet S, Plotnikoff RC, Raine K, Courneya K. Physical activity of Aboriginals with type 2 diabetes : an exploratory study. *Ethn Dis* 2005 ; 15 : 256-66.

Burton LC, Paglia MJ, German PS, et al. The effect among older persons of a general preventive visit on three health behaviors : smoking, excessive alcohol drinking, and sedentary lifestyle. *Prev Med* 1995 ; 24 : 492-7.

Burton NW, Turrell G, Oldenburg B. Participation in recreational physical activity : why do socioeconomic groups differ ? *Health Educ Behav* 2003 ; 30 : 225-44.

Calfas KJ, Long BJ, Dallis JF, et al. A controlled trial of physician counseling to promote the adoption of physical activity. *Prev Med* 1996 ; 25 : 225-33.

Calfas KJ, Long BJ, Sallis JF, et al. A controlled trail of physician counseling to promote the adoption of physical activity. *Prev Med* 1996 ; 25 : 225-33.

Canadian Fitness and Lifestyle Research Institute. Barriers to physical activity. *Progress Prev* 1996, june, bull n° 4.

Candace DR, Coleman KJ. Examining the relationships among built environment, physical activity, and body mass index in El Paso, TX. *Prev Med* 2005 ; 40 : 831-41.

Castro C, King AC, Brassington G. Telephone versus mail interventions for maintenance of physical activity in older adults. *Health Psychol* 2001 ; 20 : 438-44.

Castro C, King AC. Telephone-assisted counseling for physical activity. *Exerc Sport Sci Rev* 2002 ; 30 : 64-8.

Chen W. Chinese and American college students' motives for participation in physical activities. *Percept Mot Skills* 1998 ; 87 : 1463-70.

Cohen DA, Ashwood JS, Scott MM, et al. Public parks and physical activity among adolescent girls. *Pediatrics* 2006 ; 118 : e1381-9.

Collins R, Lee RE, Albright CL, King AC. Ready to Be Physically Active ? The Effects of a Course Preparing Low-Income Multiethnic Women to Be More Physically Active. *Health Education Behavior* 2004 ; 31 : 47-64.

Cox KL, Burke V, Gorely TJ, et al. Controlled Comparison of Retention and Adherence in Home- vs Center-Initiated Exercise Interventions in Women Ages 40-65 Years : The S.W.E.A.T. Study (Sedentary Women Exercise Adherence Trial). *Prev Med* 2003 ; 36 : 17-29.

Cox KL, Burke V, Gorely TJ, et al. Controlled comparison of retention and adherence in home- vs. center-initiated exercise interventions in women ages 40-65 years : the S.W.E.A.T. study (Sedentary Women Exercise Adherence Trial). *Prev Med* 2003 ; 36 : 17-29.

Damush TM, Stewart AL, Mill KM, et al. Prevalence and correlates of physician recommendations to exercise among older adults. *J Gerontol A Biol Sci Med Sci* 1999 ; 54 : M423-7.

de Andrade Bastos A, Salguero A, Gonzalez-Boto R, Marquez S. Motives for participation in physical activity by Brazilian adults. *Percept Mot Skills* 2006 ; 102 : 358-67.

De Bourdeaudhuij I, Sallis JF. Relative contribution of psychosocial variables to the explanation of physical activity in three population-based adult samples. *Prev Med* 2002 ; 34 : 279-88.

Deforche BI, De Bourdeaudhuij IM, Tanghe AP. Attitude toward physical activity in normal-weight, overweight and obese adolescents. *J Adolesc Health* 2006 ; 38 : 560-8.

Delahanty LM, Conroy MB, Nathan DM, Diabetes Prevention Program Research Group. Psychological predictors of physical activity in the diabetes prevention program. *J Am Diet Assoc* 2006 ; 106 : 698-705.

Desmond AW, Conrad KM, Montgomery A, Simon KA. Factors associated with male workers' engagement in physical activity. *AAOHN J* 1993 ; 41 : 73-83.

Dishman RK, Buckworth J. Increasing physical activity : a quantitative synthesis. *Med Sci Sports Exerc* 1996 ; 28 : 706-19

Dishman RK, Motl RW, Saunders R, et al. Self-efficacy partially mediates the effect of a school-based physical-activity intervention among adolescent girls. *Prev Med* 2004 ; 38 : 628-36.

Dishman RK, Oldenburg B, O'Neal H, Shephard RJ. Worksite physical activity interventions. *Am J Prev Med* 1998 ; 15 : 344-61.

Dishman RK. Compliance/adherence in health-related exercise. *Health Psychol* 1982 ; 3 : 237-67.

Dishman RK. *Exercise adherence : its impact on public health*. Champaign, IL : Human Kinetics, 1988 : 1-9.

Dowell AC, Ochera JJ, Hilton SR, et al. Prevention in practice : results of a 2-year follow-up of routine health promotion interventions in general practice. *Fam Pract* 1996 ; 13 : 357-62.

Dunn AL, Marcus BH, Kampert JB, et al. Comparison of lifestyle and structured interventions to increase physical activity and cardiorespiratory fitness : a randomised trial. *JAMA* 1999 ; 281 : 327-34.

Dwyer JJ, Allison KR, Goldenberg ER, et al. Adolescent girls' perceived barriers to participation in physical activity. *Adolescence* 2006 ; 41 : 75-89.

Eakin EG, Brown WJ, Marshall AL, et al. Physical activity promotion in primary care : bridging the gap between research and practice. *Am J Prev Med* 2004 ; 27 : 297-303.

Eaton CB, Menard LM. A systematic review of physical activity promotion in primary care office settings. *Br J Sports Med* 1998 ; 32 : 11-6.

Elley CR, Kerse N, Arroll B, Robinson E. Effectiveness of counselling patients on physical activity in general practice : cluster randomised controlled trial. *Br Med J* 2003 ; 326 : 793.

Ewing ME, Seefeldt V. *Participation and attrition patterns in American-agency-sponsored and interscholastic sports : An executive summary*. Final report. North Palm Beach, Sporting Goods Manufacturer's Assoc., 1989.

Faulkner G, Biddle S. Predicting physical activity promotion in health care settings. *Am J Health Promot* 2001 ; 16 : 98-106.

Fischbein M, Ajzen I. *Belief, attitude, intention and behavior : an introduction to theory and research*. Reading MA : Addison-Wesley, 1975

Frederick CM, Ryan RM. Differences in motivation for sport and exercise and their relationships with participation and mental health. *J Sport Behavior* 1993 ; 16 : 125-45.

Galuska DA, Fulton JE, Powell KE, et al. Pediatrician Counseling About Preventive Health Topics : Results From the Physicians' Practices Survey, 1998 -1999. *Pediatrics* 2002 ; 109 : e83.

Garet M, Chamoux A, Margot AC, Brunet F. Évaluation de l'activité physique et de la tolérance à l'effort chez les personnes lourdement handicapées vivant en institution. *Rev Eur Handicap Mental* 2004 ; 29 : 19-35.

Genkinger JM, Jehn ML, Sapun M, et al. Does weight status influence perceptions of physical activity barriers among African-American women ? *Ethn Dis* 2006 ; 16 : 78-84.

Godin G, Valois P, Jobin J, Ross A. prediction of intention to exercise of individuals who have suffered from coronary heart disease. *J Clin Psychol* 1991 ; 47 : 762-72.

Godin G, Valois P, Lepage L. The pattern of influence of perceived behavioral control upon exercising behavior : an application of Ajzen's theory of planned behavior. *J Behav Med* 1993 ; 16 : 81-102.

Grant AM, Niyonsenga T, Dion I, et al. Cardiovascular disease. Physicians attitudes toward prevention and treatment. *Can Fam Physician* 1998 ; 44 : 780-7.

Halbert JA, Silagy CA, Finucane P, et al. Physical activity and cardiovascular risk factors : effect of advice from an exercise specialist in Australian general practice. *Med J Aust* 2000 ; 173 : 84-7.

Halbert JA, Silagy CA, Finucane P, et al. Recruitment of older adults for a randomized, controlled trial of exercise advice in a general practice setting. *J Am Geriatr Soc* 1999 ; 47 : 477-81.

Hallam JS, Petosa R. The long-term impact of a four-session work-site intervention on selected social cognitive theory variables linked to adult exercise adherence. *Health Educ Behav* 2004 ; 31 : 88-100.

Harland J, White M, Drinkwater C, et al. The Newcastle exercise project : a randomised controlled trial of methods to promote physical activity in primary care. *Br Med J* 1999 ; 319 : 828-32.

Hillsdon M, Thorogood M, White I, Foster C. Advising people to take more exercise is ineffective : a randomized controlled trial of physical activity promotion in primary care. *Int J Epidemiol* 2002 ; 31 : 808-15.

Hirvensalo M, Heikkinen E, Lintunen T, Rantanen T. Recommendations for and warnings against physical activity given to older people by health care professionals. *Prev Med* 2005 ; 41 : 342-7.

Hochbaum GM. *Public participation in medical screening programs. A sociopsychological study.* US Public Service Publication n° 572. Washington DC : US Gov Print Office, 1958.

Hofstetter CR, Hovell MF, Macera C, et al. Illness, injury, and correlates of aerobic exercise and walking : a community study. *Res Q Exerc Sport* 1991 ; 62 : 1-9.

Hume C, Ball K, Salmon J. Development and reliability of a self-report questionnaire to examine children's perceptions of the physical activity environment at home and in the neighbourhood. *Int J Behav Nutr Phys Act* 2006 ; 3 : 16.

Humpel N, Marshall AL, Iverson D, et al. Trial of print and telephone delivered interventions to influence walking. *Prev Med* 2004 ; 39 : 635-41.

Imperial Cancer Research fund OXCHECK study group. Effectiveness of health checks conducted by nurses in primary care : final results of the OXCHECK study. *Br Med J* 1995 ; 310 : 1099-104.

Jackson NW, Howes FS, Gupta S, et al. Interventions implemented through sporting organisations for increasing participation in sport. *The Cochrane Database of Systematic Reviews* 2005, Issue 2. Art. No. : CD004812.pub2. DOI : 10.1002/14651858.CD004812.pub2.

Kaewthummanukul T, Brown KC. Determinants of employee participation in physical activity : critical review of the literature. *AAOHN J* 2006 ; 54 : 249-61.

Kelly RB. Controlled trial of a time-efficient method of health promotion. *Am J Prev Med* 1988 ; 4 : 200-7.

Kerse NM, Flicker L, Jolley D, et al. Improving the health behaviours of elderly people : randomised controlled trial of a general practice education programme. *Br Med J* 1999 ; 319 : 683-7.

King AC, Haskell WL, Taylor CB, et al. Group vs home-based exercise training in healthy older men and women : a community-based trial. *JAMA* 1991 ; 266 : 1535-42.

King AC, Haskell WL, Young DR, et al. Long-term effects of varying intensities and formats of physical activity on participation rates, fitness, and lipoproteins in men and women aged 50 – 65 years. *Circulation* 1995 ; 91 : 2596-604.

King AC, Rejeski JW, Buchner DM. Physical activity interventions targeting older adults. A critical review. *Am J Prev Med* 1998 ; 15 : 316-33.

King AC, Sallis JF, Dunn AL, et al. Overview of the Activity Counseling Trial (ACT) intervention for promoting physical activity in primary health care settings. Activity Counseling Trial Research Group. *Med Sci Sports Exerc* 1998 ; 30 : 1086-96.

Kirk AF, Higgins LA, Hughes AR, et al. A randomized, controlled trial to study the effect of exercise consultation on the promotion of physical activity in people with type 2 diabetes : a pilot study. *Diabet Med* 2001 ; 18 : 877-82.

Lamb SE, Bartlett HP, Ashley A, Bird W. Can lay-led walking programmes increase physical activity in middle aged adults ? A randomised controlled trial. *J Epidemiol Comm Health* 2002 ; 56 : 246-52.

Lasheras L, Aznar S, Merino B, Lopez EG. Factors associated with physical activity among Spanish Youth through the National Health Survey. *Prev Med* 2001 ; 32 : 455-64.

Laukkanen RM, Kalaja MK, Kalaja SP, et al. Heart rate during aerobics classes in women with different previous experience of aerobics. *Eur J Appl Physiol* 2001 ; 84 : 64-8.

Lawlor DA, Keen S, Neal RD. Increasing population levels of physical activity through primary care : GPs' knowledge, attitudes and self-reported practice. *Fam Pract* 1999 ; 16 : 250-4.

Les adolescents et le sport. Enquête 2001. Paris : Ministère de la Jeunesse, des Sports et de la Vie associative et INSEP, 2004.

Les pratiques sportives en France. Enquête 2000. Paris : Ministère des sports et INSEP, 2002.

Leslie E, Johnson-Kozlow M, Sallis JF, et al. Reliability of moderate-intensity and vigorous physical activity stage of change measures for young adults. *Prev Med* 2003 ; 37 : 173-81.

Lewis BL, Lynch WD. The effect of physician advice on exercise behavior. *Prev Med* 1993 ; 22 : 110-21.

Lim K, Taylor L. Factors associated with physical activity among older people– a population-based study. *Prev Med* 2005 ; 40 : 33-40.

Lindwall M, Martin Ginis KA. Moving towards a favorable image : the self-presentational benefits of exercise and physical activity. *Scand J Psychol* 2006 ; 47 : 209-17.

Lippke S, Ziegelmann JP, Schwarzer R. Stage-specific adoption and maintenance of physical activity : testing a three-stage model. *Psychol Sport Exerc* 2005 ; 6 : 585-603.

Lodgsdon DN, Lazaro CM, Meier RV. The feasibility of behavioral risk reduction in primary medical care. *Am J Prev Med* 1989 ; 5 : 249-56.

Lombard DN, Lombard TN, Winett RA. Walking to meet health guidelines : the effect of prompting frequency and prompt structure. *Health Psychol* 1995 ; 14 : 164-70.

London Sports Forum for Disabled People. *Sport achieving equality.* Annual Review 2003-2004. London, 2004.

Maloof RM, Zabik RM, Dawson ML. The effect of use of a personal trainer on improvement of health related fitness for adults. *Med Sci Sports Exerc* 2001 ; 33 : s74.

Marcus BH, Emmons K, Simkin-Silverman L, et al. Evaluation of motivationally tailored vs. standard self-help physical activity interventions. *Am J Health Promot* 1998 ; 12 : 246-53.

Marshall AL, Bauman A, Owen N. Reaching out to promote physical activity : a statewide randomized controlled trial of a stage-targeted intervention. *Am J Health Promot* 2004 ; 18 : 283-7.

Marshall SJ, Biddle SJ. The transtheoretical model of behavior change : a meta-analysis of applications to physical activity and exercise. *Ann Behav Med* 2001 ; 4 : 229-46.

Marti B, Seleger U, Schwyn C, Denoth J. Physiological, social and motivational characteristics of fitness center clients. *Schweiz Z Sportmed* 1989 ; 37 : 233-9.

McAuley E, Blissmer B. Self-efficacy determinants and consequences of physical activity. *Exerc Sport Sci Rev* 2000 ; 28 : 85-8.

McAuley E. Self-efficacy and the maintenance of exercise participation in older adults. *J Behav Med* 1993 ; 16 : 103-13.

McClaran SR. The effectiveness of personal training on changing attitudes towards physical activity. *J Sports Sci Med* 2003 ; 2 : 10-4.

McKenna J, Henderson L, Baic S. A survey to assess physical activity promotion by Registered Dietitians. *J Hum Nutr Dietet* 2004 ; 17 : 63-9.

McKenna J, Naylor PJ, McDowell N. Barriers to physical activity promotion by general practitioners and practice nurses. *Br J Sports Med* 1998 ; 32 : 242-7.

Merrill RM, Thygeson AL. Religious Preference, Church Activity, and Physical Exercise. *Prev Med* 2001 ; 33 : 38-45.

Mota J, Almeida M, Santos P, Ribeiro JC. Perceived Neighborhood Environments and physical activity in adolescents. *Prev Med* 2005 ; 41 : 834-6.

Navuluri RB. Gender differences in the factors related to physical activity among adults with diabetes. *Nursing Health Sci* 2000 ; 2 : 191-99.

Nishida Y, Suzuki H, Wang DH, Kira S. Psychological determinants of physical activity in Japanese female employees. *J Occup Health* 2003 ; 45 : 15-22.

Norris SL, Grothaus MA, Buchner DM, Pratt M. Effectiveness of physician-based assessment and counseling for exercise in a staff model HMO. *Prev Med* 2000 ; 30 : 513-23.

Nygaard I, Girts T, Pharm D, et al. Is Urinary Incontinence a Barrier to Exercise in Women ? *Obstetrics Gynecol* 2005 ; 106 : 307-14.

O'Brien Cousins S, Gillis MM. « Just do it... before you talk yourself out of it » : the self-talk of adults thinking about physical activity. *Psychol Sport Exerc* 2005 ; 6 : 313-34.

O'Neill K, Reid G. Perceived barriers to physical activity by older adults. *Can J Public Health* 1991 ; 82 : 392-6.

Ortega-Sanchez R, Jimenez-Mena C, Cordoba-Garcia R, et al. The effect of office-based physician's advice on adolescent exercise behavior. *Prev Med* 2004 ; 38 : 219-26.

Palmer T, Jaworski CA. Exercise prescription for underprivileged minorities. *Curr Sports Med Rep* 2004 ; 3 : 344-8.

Patureau F. *Les pratiques culturelles des jeunes*. Paris : La Documentation française, 1992.

Perri MG, Anton SD, Durning PE, et al. Adherence to exercise prescriptions : effects of prescribing moderate versus higher levels of intensity and frequency. *Health Psychol* 2002 ; 21 : 452-8.

Petosa RL, Hartz BV, Cardina CE, Suminski RR. Social cognitive theory variables associated with physical activity among high school students. *Int J Sports Med* 2005 ; 26 : 158-63.

Pinto BM, Friedman R, Marcus BH, et al. Effects of a computer-based, telephone-counseling system on physical activity. *Am J Prev Med* 2002 ; 23 : 113-20.

Pinto BM, Goldstein MG, DePue JD, Milan FB. Acceptability and feasibility of physician-based activity counseling. The PAL project. *Am J Prev Med* 1998 ; 15 : 95-102.

Pinto BM, Lynn H, Marcus BH, et al. Physician-based activity counseling : intervention effects on mediators of motivational readiness for physical activity. *Ann Behav Med* 2001 ; 23 : 2-10.

Plotnikoff RC, Mayhew A, Birkett N, et al. Age, gender, and urban-rural differences in the correlates of physical activity. *Prev Med* 2004 ; 39 : 1115-25.

Pohjonen T, Ranta R. Effects of Worksite Physical Exercise Intervention on Physical Fitness, Perceived Health Status, and Work Ability among Home Care Workers : Five-Year Follow-up. *Prev Med* 2001 ; 32 : 465-75.

Poulsen AA, Ziviani J.-M. Health enhancing physical activity : Factors influencing engagement patterns in children. *Aust Occupational Therapy J* 2004 ; 51 : 69-79.

Prochaska JO, DiClemente CC, Norcross JC. In search of how people change. Applications to addictive behaviors. *Am Psychol* 1992 ; 47 : 1102-14.

Puig Ribera A, McKenna J, Riddoch C. Attitudes and practices of physicians and nurses regarding physical activity promotion in the Catalan primary health-care system. *Eur J Public Health* 2005 ; 15 : 569-75.

Reis JP, Bowles HR, Ainsworth BE, et al. Nonoccupational Physical Activity by Degree of Urbanization and U.S. Geographic Region. *Med Sci Sports Exerc* 2004 ; 36 : 1093-8.

Rejeski WJ, Brawley LR. Shaping active lifestyles in older adults : a group-facilitated behaviour change intervention. *Ann Behav Med* 1997 ; 19 : S106

Reynolds KD, Killen JD, Bryson SW, et al. Psychological predictors of physical activity in adolescents. *Prev Med* 1990 ; 19 : 541-51.

Rhodes RE, Brown SG, McIntyre CA. Integrating the perceived neighborhood environment and the theory of planned behavior when predicting walking in a Canadian adult sample. *Am J Health Promot* 2006 ; 21 : 110-8.

Rimmer JH, Riley B, Wang E, et al. Physical activity participation among persons with disabilities : barriers and facilitators. *Am J Prev Med* 2004 ; 26 : 419-25.

Rimmer JH, Rubin SS, Braddock D. Barriers to exercise in African American women with physical disabilities. *Arch Phys Med Rehabil* 2000 ; 81 : 182-8.

Rimmer JH. The conspicuous absence of people with disabilities in public fitness and recreation facilities : lack of interest or lack of access ? *Am J Health Promot* 2005 ; 19 : 327-9.

Rogers LQ, Shah P, Dunnington G, et al. Social cognitive theory and physical activity during breast cancer treatment. *Oncol Nurs Forum* 2005 ; 32 : 807-15.

Romero AJ, Robinson TN, Kraemer HC, et al. Are Perceived Neighborhood Hazards a Barrier to Physical Activity in Children ? *Arch Pediatr Adolesc Med* 2001 ; 155 : 1143-8.

Romero AJ. Low-income neighborhood barriers and resources for adolescents' physical activity. *J Adolesc Health* 2005 ; 36 : 253-9.

Rütten A, Abu-Omar K. Perceptions of environmental opportunities for physical activity in the European Union. *Soz Präventivmed* 2004 ; 49 : 310-7.

Ryan RM, Frederick CM, Lepes D, et al. Intrinsic motivation and exercise adherence. *Int J Sport Psychol* 1997 ; 28 : 335-54.

Sallis J F, Hovell MF, Hofstetter CR, et al. A multivariate study of determinants of vigorous exercise in a community sample. *Prev Med* 1989 ; 18 : 20-34

- Sallis JF, Haskell WL, Fortmann SP, et al. Predictors of adoption and maintenance of physical activity in a community sample. *Prev Med* 1986 ; 15 : 331-41.
- Schofield L, Mummery WK, Schofield G. Effects of a Controlled Pedometer-Intervention Trial for Low-Active Adolescent Girls. *Med Sci Sports Exerc* 2005 ; 37 : 1414-20.
- Schonberg MA, Marcantonio ER, Wee CC. Receipt of exercise counseling by older women. *J Am Geriatr Soc* 2006 ; 54 : 619-26.
- Schutzer KA, Graves BS. Barriers and motivations to exercise in older adults. *Prev Med* 2004 ; 39 : 1056-61.
- Sciamanna CN, Goldstein MG, Marcus BH, et al. Accuracy of recall of exercise counseling among primary care patients. *Prev Med* 2004 ; 39 : 1063-7.
- Sechrist KR, Walker SN, Pender NJ. Development and psychometric evaluation of the Exercise Benefits/Barriers Scale. *Res Nurs Health* 1987 ; 10 : 357-365.
- Shephard RJ. Worksite fitness and exercise programs : a review of methodology and health impact. *Am J Health Promot* 1996 ; 10 : 436-52.
- Sherman SE, Hershman WY. Exercise counselling : how do general internists do ? *J Gen Intern Med* 1993 ; 8 : 243-8.
- Silberstein LR, Striegel-Moore RH, Timko C, Rodin J. Behavioral and psychological implications of body dissatisfaction : do men and women differ ? *Sex Roles* 1988 ; 19 : 219-32.
- Simons-Morton DG, Hogan P, Dunn AL, Pruitt L, King AC, Levine Bd, et al. Characteristics of inactive primary care patients : baseline data from the Activity Counseling Trial. *Prev Med* 2000 ; 31 : 513-21.
- Smith BJ, Bauman AE, Bull FC, et al. Promoting physical activity in general practice : a controlled trial of written advice and information materials. *Br J Sports Medicine* 2000 ; 34 : 262-7.
- Snodgrass SJ, Rivett DA, Mackenzie LA. Perceptions of older people about falls injury prevention and physical activity. *Australasian J Ageing* 2005 ; 24 : 114-8.
- Sorensen JB, Skovgaard T, Puggaard L. Exercise on prescription in general practice : a systematic review. *Scand J Prim Health Care* 2006 ; 24 : 69-74.
- Stach-Lempinen B, Nygard CH, Laippala P, et al. Is physical activity influenced by urinary incontinence ? *Int J Obst Gynaecol* 2004 ; 111 : 475-80.
- Swinburn B, Walter L, Arroll B, et al. The Green Prescription Study : a randomized controlled trial of exercise prescription in general practice. *Am J Public Health* 1998 ; 88 : 288-91.
- Taggart HM, Connor SE. The relation of exercise habits to health beliefs and knowledge about osteoporosis. *J Am College Health* 1995 ; 44 : 127-30.
- Tan D, Zwar NA, Dennis SM, Vagholkar S. Weight management in general practice : what do patients want ? *Med J Aust* 2006 ; 185 : 73-5.
- The Writing Group for the Activity Counseling Trial Research Group. Effects of physical activity counseling in primary care : the Activity Counseling Trial : a randomized controlled trial. *JAMA* 2001 ; 286 : 677-87.
- Thomas RK, Fulton JE, Galuska DA. Weight counseling patterns of U.S. pediatricians. *Obes Res* 2004 ; 12 : 161-9.
- Thorpe RJ Jr, Simonsick EM, Brach JS, et al. Dog ownership, walking behavior, and maintained mobility in late life. *J Am Geriatr Soc* 2006 ; 54 : 1419-24.
- Timperio A, Crawford D, Telford A, Salmon J. Perceptions about the local neighborhood and walking and cycling among children. *Prev Med* 2004 ; 38 : 39-47.
- Titze S, Stronegger W, Owen N. Prospective study of individual, social, and environmental predictors of physical activity : women's leisure running. *Psycho Sport Exerc* 2005 ; 6 : 363-76.
- Trost SG, Owen N, Bauman AE, Sallis JF, Brown W. Correlates of adults' participation in physical activity : review and update. *Med Sci Sports Exerc* 2002 ; 12 : 1996-2001.
- Van Sluijs EM, Van Poppel MN, Twisk JW, et al. The positive effect on determinants of physical activity of a tailored, general practice-based physical activity intervention. *Health Educ Res* 2005 ; 20 : 345-56.
- Verheijden MW, Bakx JC, Delemarre IC, et al. GPs' assessment of patients' readiness to change diet, activity and smoking. *Br J Gen Pract* 2005 ; 55 : 452-7.
- Walseth K, Fasting K. Islam's view of physical activity and sport. *Int Rev Soc Sport* 2003 ; 38 : 45-60.

- Warms CA, Belza BL, Whitney JD, et al. Lifestyle physical activity for individuals with spinal cord injury : a pilot study. *Am J Health Promot* 2004 ; 18 : 288-91.
- Wee CC, McCarthy EP, Davis RB, Philips RS. Physical counseling about exercise. *JAMA* 1999 ; 282 : 1583-8.
- Wilcox BS, Castro C, King AC, et al. Determinants of leisure time physical activity in rural compared with urban older and ethnically diverse women in the United States. *J Epidemiol Community Health* 2000 ; 54 : 667-72.
- Williams DM, Anderson ES, Winett RA. A review of the outcome expectancy construct in physical activity research. *Ann Behav Med* 2005 ; 29 : 70-9.
- Wing RR, Jeffrey RW, Pronk N, Hellerstedt WL. Effects of a personal trainer and financial incentives on exercise adherence in overweight women in a behavioral weight loss program. *Obesity Research* 1996 ; 5 : 457-62.
- World Health Organization. *World Health Report 2001*. Geneva, WHO, 2001
- Yordy GA, Lent RW. Predicting aerobic exercise participation : social cognitive, reasoned action, and planned behavior models. *J Sport Exerc Psychol* 1993 ; 15 : 363-74.
- Zunft HJF, Friebe D, Seppelt B, et al. Perceived benefits and barriers to physical activity in a nationally representative sample in the European Union. *Pub Health Nutr* 1999 ; 2 : 153-160.

STRATÉGIES DE PROMOTION DE LA SANTÉ

- Active Start : A Statement of Physical Activity Guidelines for Children Birth to Five Years. National Association for Sport & Physical Education. Oxon Hill, AAHPERD : 2002.
- American College of Sports Medicine Position Stand. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Med Sci Sports Exerc.* 1998 ; 6 : 975-991.
- American College of Sports Medicine. Position stand on the recommended quantity and quality of exercise for developing and maintaining fitness in healthy adults. *Med Sci Sports* 1978 ; 10 : 7-10.
- American College of Sports Medicine. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness in healthy adults. *Med Sci Sports Exerc* 1990 ; 22 : 265-274.
- Anhøj J, Holm Jensen A. Using the Internet for Life Style Changes in Diet and Physical Activity : A Feasibility Study. *J Med Internet Res* 2004 ; 6 (3) : e28.
- Bauman AE, Bellew B, Owen N, Vita P. Impact of an Australian mass media campaign targeting physical activity in 1998. *Am J Prev Med* 2001 ; 21 : 41-7.
- Biddle S, Sallis J, Cavill N (Eds). *Young and active ? Young people and health enhancing physical activity – evidence and implications*. Health Education Authority. London, England, 1998
- Blair SN. 1993 CH McCloy Research Lecture : Physical activity, physical fitness, and health. *Res Quart Exerc Sport* 1993 ; 64 : 365-376.
- Burdette HL, Whitaker RC. Resurrecting free play in young children : looking beyond fitness and fatness to attention, affiliation, and affect. *Arch Pediatr Adolesc Med* 2005 ; 159 : 46-50.
- Cavill N, Biddle S, Sallis JF. Health enhancing physical activity for young people : Statement of the United Kingdom Expert Consensus Conference. *Pediatr Exerc Sci.* 2001 ; 13 : 12-25.
- Centers for Disease Control and Prevention. Increasing physical activity : a report on recommendations of the Task Force on Community Preventive Services. *MMWR* 2001 ; 50 (No. RR-18).
- Connolly P, McKenzie TL. Effects of a games intervention on the physical activity levels of children at recess. *Res Q Exerc Sport* 1995 ; 66 : A60.
- Corti BG, Donovan RJ. The relative influence of individual, social and physical environment determinants of physical activity. *Soc Sci Med* 2002 ; 54 : 1793, 1812.
- Cox M, Schofield G, Greasley N, Kolt GS. Pedometer steps in primary school-aged children : a comparison of school-based and out-of-school activity. *J Sci Med Sport* 2006 ; 9 : 91-7.
- Craig CL, Cragg SE, Tudor-Locke C, Bauman A. Impact proximal de Canada en mouvement : relation entre la notoriété de la campagne et la propriété et l'utilisation d'un podomètre. *Can J Public Health* 2006 ; 97 : S22-9.
- Epstein L, Paluch R, Kalakanis L, et al. How much activity do youth get ? A quantitative review of heart -rate measured activity. *Pediatrics* 2001 ; 108 (3) [http : //www.pediatrics.org/cgi/content/full/3/e44](http://www.pediatrics.org/cgi/content/full/3/e44).

- Ernst MP, Pangrazi RP. Effects of a physical activity program on children's activity levels and attraction to physical activity. *Pediatr Exerc Sci* 1999 ; 11 : 393-405.
- Exercise for Health.WHO/FIMS Committee on Physical Activity for Health. *Bull WHO* 1995 ; 2 : 135-136.
- Faulkner G, Finlay SJ. Canada en mouvement : analyse intensive des reportages des médias, de la création à la réception de la campagne. *Can J Public Health* 2006 ; 97 : S17-21.
- Finlay SJ, Faulkner G. Physical activity promotion through the mass media : Inception, production, transmission and consumption. *Prev Med* 2005 ; 40 : 121-30.
- Gates D, Brehm B, Hutton S, et al. Changing the work environment to promote wellness : a focus group study. *AAOHN J* 2006 ; 54 : 515-20.
- Hill JW. Exercise prescription. *Prim Care* 1987 ; 14 : 817-825
- Hillsdon M, Cavill N, Nanchahal K, et al. National level promotion of physical activity : results from England's ACTIVE for LIFE campaign. *J Epidemiol Community Health* 2001 ; 55 : 755-61.
- Increasing Physical Activity through Community design. A guide for Public health practitioners.* Washington : National Center for Bicycling and Walking, may 2002.
- Irwin JD, He M, Bouck LM, Tucker P, Pollett GL. Preschoolers' physical activity behaviours : parents' perspectives. *Can J Public Health* 2005 ; 96 : 299-303.
- Jago R, Baranowski T. Non-curricular approaches for increasing physical activity in youth : a review. *Prev Med* 2004 ; 39 : 157-63.
- Kahn EB, Ramsey LT, Brownson RC, et al. The effectiveness of interventions to increase physical activity. A systematic review. *Am J Prev Med* 2002 ; 22 : S73-107.
- La Déclaration de Jakarta sur la promotion de la santé au XXIe siècle.* HPR/HEP/41CHP/BR/97.4.OMS, Genève, 1997.
- Leslie E, Marshall AL, Owen N, Bauman A. Engagement and retention of participants in a physical activity website. *Prev Med* 2005 ; 40 : 54-9.
- Marcus BH, Owen N, Forsyth LH, et al. Physical activity interventions using mass media, print media, and information technology. *Am J Prev Med* 1998 ; 15 : 362-78.
- Marks JT, Campbell MK, Ward DS, et al. A comparison of Web and print media for physical activity promotion among adolescent girls. *J Adolesc Health* 2006 ; 39 : 96-104.
- McKay HG, King D, Eakin EG, et al. The Diabetes Network internet-based physical activity intervention. *Diabetes Care* 2001 ; 24 : 1328-34.
- Merom D, Bauman A, Vita P, Close G. An environmental intervention to promote walking and cycling — the impact of a newly constructed Rail Trail in Western Sydney. *Prev Med* 2003 ; 36 : 235-42.
- Miles A, Rapoport L, Wardle J, et al. Using the mass-media to target obesity : an analysis of the characteristics and reported behavior change of participants in the BBC's 'Fighting Fat, Fighting Fit' campaign. *Health Educ Res* 2001 ; 16 : 357-72
- Napolitano MA, Fotheringham M, Tate D, et al. Evaluation of an Internet-based physical activity intervention : a preliminary investigation. *Ann Behav Med* 2003 ; 25 : 92-9.
- Nigg CR. Technology's influence on physical activity and exercise science : the present and the future. *Psychol Sport Exerc* 2003 ; 4 : 57-65.
- Obesity Prevention for Health Care Systems—Executive Summary and Literature Review,* California Obesity Prevention Initiative Health Systems Workgroup, California Department of Health Services, 2002.
- O'Hara Tompkins N, Zizzi S, et al. School-based opportunities for physical activity in West Virginia public schools. *Prev Med* 2004 ; 39 : 834-40.
- Pate RR, Pratt M, Blair SN, et al. Physical activity and public health : a recommendation from the Center for Disease Control and Prevention and the ACSM. *J Am Med Assoc* 1995 ; 273 : 402-407.
- Pate RR, Trost SG, Mullis R, et al. Community Interventions to Promote Proper Nutrition and Physical Activity among Youth. *Prev Med* 2000 ; 31 : S138-69.
- Poscente N, Rothstein M, Irvine J. Using marketing research methods to evaluate a stage-specific intervention. *Am J Health Behav* 2002 ; 26 : 243-51.
- Recommendations to Increase Physical Activity in Communities. *Am J Prev Med* 2002 ; 22 : S67-72.
- Reger B, Cooper L, Booth-Butterfield S, et al. Wheeling Walks : a community campaign using paid media to encourage walking among sedentary older adults. *Prev Med* 2002 ; 35 : 285-92.

- Richter KP, Harris KJ, Paine-Andrews A, et al. Measuring the Health Environment for Physical Activity and Nutrition among Youth : A Review of the Literature and Applications for Community Initiatives. *Prev Med* 2000 ; 31 : S98-111.
- Robbins LB, Gretebeck KA, Kazanis AS, Pender NJ. Girls on the move program to increase physical activity participation. *Nurs Res* 2006 ; 55 : 206-16.
- Rowland D, Diguseppi C, Gross M, et al. Randomised controlled trial of site specific advice on school travel patterns. *Arch Dis Child* 2003 ; 88 : 8-11.
- Sallis J, Patrick K. Physical activity guidelines for adolescents : Consensus statement. *Pediatr Exerc Sci* 1994 ; 6 : 302-14
- Sallis JF, Johnson MF, Calfas KJ, et al. Assessing perceived physical environmental variables that may influence physical activity. *Res Q Exerc Sport* 1997 ; 68 : 345 -51.
- Sallis JF, Nader PR, Broyles SL, et al. Correlates of physical activity at home in Mexican-American and Anglo-American preschool children. *Health Psychol* 1993 ; 12 : 390-8.
- Sciamanna CN, Lewis B, Tate D, Napolitano MA, et al. User attitudes toward a physical activity promotion website. *Prev Med* 2002 ; 35 : 612-5.
- Simon C, Wagner A, DiVita C, et al. Intervention centred on adolescents' physical activity and sedentary behaviour (ICAPS) : concept and 6-month results. *Int J Obes Relat Metab Disord* 2004 ; 28 : S96-103.
- Simon C, Wagner A, Platat C, et al. ICAPS : a multilevel program to improve physical activity in adolescents. *Diabetes Metab* 2006 ; 32 : 41-9.
- Stahl T, Rütten A, Nutbeam D, et al. The importance of the social environment for physical activity lifestyle – results from an international study. *Soc Sci Med* 2001 ; 52 : 1-10.
- Stahl T, Rütten A, Nutbeam D, Kannas L. The importance of policy orientation and environment on physical activity participation- a comparative analysis between Eastern Germany, Western Germany and Finland. *Health Promotion Int* 2002 ; 17 : 235-46.
- Strategic direction and advice for increasing safe participation in physical activity in the Australian Defence Force : A report to the Defence Health Service.* Victoria : Monash University Accident Research Centre, 2002.
- Stratton G. Promoting children's physical activity in primary schools : an intervention study using playgrounds markings. *Ergonomics* 2000 ; 43 : 1538-46.
- Tate AK, Petruzzello SJ. Varying the intensity of acute exercise : implication for changes in affect. *J Sports Med Phys Fitness* 1995 ; 35 : 1-8.
- The Heidelberg guidelines for promoting physical activity among older persons.* Ageing and Health Programme. Geneva, WHO, 1996.
- The promotion of physical activity in Scotland : strategic statement.* Edimbourg, Health Education Board for Scotland : 2003.
- Troped PJ, Saunders RP, Pate RR, et al. Association between self-reported and objective physical environmental factors and use of community Rail-Trail. *Prev Med* 2001 ; 32 : 191-200.
- US Department of Health and Human Services. *Physical activity evaluation handbook.* Atlanta : Centers for Disease Control and Prevention, 2002.
- Van Beurden E, Barnett LM, Zask A, et al. Can we skill and activate children through primary school physical education lessons ? "Move it Groove it" – a collaborative health promotion intervention. *Prev Med* 2003 ; 36 : 493-501.
- Wallack L. Mass media campaigns : the odds against finding behavior change. *Health Educ Q* 1981 ; 8 : 209-60.
- Warburton DE, Nicol CW, Bredin SS. Prescribing exercise as preventive therapy. *Can Med Assoc J* 2006 ; 174 : 961-74.
- WHO. *Health Promotion Glossary.* WHO/HPR/HEP/98.1. Geneva : WHO, 1999.
- WHO/CDC Partners Meeting on Implementation of WHO Physical Activity Strategy,* Miami 13-15 Dec 2004. Action Plan February 2005. WHO/NMH/CHP/HPR-PCD/PAH/hba.



Éléments de conseil en activité physique

Thomas S, Reading J, Shephard RJ. Revision of the Physical Activity Readiness Questionnaire (PAR-Q). *Can J Sport Sci* 1992 ; 17 : 338-45.